

BURNOUT, RESILIENCE AND LIFE SATISFACTION AMONG DOCTORS WORKING WITH HIGH-RISK PATIENTS

Ms. Maha Iqbal*¹

¹Department of Psychiatry and Behavioral Sciences, Faisalabad Medical University

¹ansarimahaiqbal@gmail.com

Mr. Gulam Mustafa Niazi²

²Department of Psychiatry and Behavioral Sciences, Faisalabad Medical University

²ghulammustfakhanniazi80@gmail.com

Prof. Dr. Imtiaz Ahmad Dogar³

³Department of Psychiatry and Behavioral Sciences, Faisalabad Medical University

³psycon2005fsd@yahoo.com

Ms. Sitara Ashraf⁴

⁴Department of Psychiatry and Behavioral Sciences, Faisalabad Medical University

⁴sitaraashraf4@gmail.com

Author Details

Keywords:

Burnout, Resilience, Life satisfaction, Doctors, High Risk Patients, Medicine and Surgery Department

Received on 03 March, 2026

Accepted on 17 May, 2026

Published on 06 Jun, 2026

Corresponding E-mails & Authors*:

Ms. Maha Iqbal

ansarimahaiqbal@gmail.com

Abstract

Burnout has become a major problem among doctors that negatively affects not only the individual and the patients, but also the healthcare systems. Burned-out doctors show an increased likelihood of generating errors and poor decision-making. Doctors who possess resilience, along with life satisfaction, are less prone to stressors, thus preventing them from burnout. This study aimed to analyze the correlations between burnout, resilience, and life satisfaction among doctors (N = 151, Mage = 26.32; SD = 2.864) dealing with high-risk patients in the Medicine and Surgery

departments of Allied Hospitals 1 and 2, Faisalabad, Pakistan. The data were collected using validated self-report questionnaires, including the Copenhagen Burnout Inventory (CBI), the Brief Resilience Scale, and the Satisfaction with Life Scale. Data was analyzed

DOI: <http://doi.org/10.5281/zenodo.20569901>

using both the Pearson Correlation Coefficient and hierarchical regression techniques. The results reveal a positive and significant relationship between resilience and life satisfaction ($r = 0.42$, $p < 0.001$), while it also reports a significant negative relationship between resilience and burnout ($r = -0.18$, $p < 0.05$). Similarly, results also showed a negative association between burnout and life satisfaction ($r = -0.36$, $p < 0.001$). Overall results also showed that both resilience ($\beta = 0.39$, $p < 0.001$) and personal burnout ($\beta = -0.32$, $p < 0.001$) significantly predicted life satisfaction. These findings have implications for doctors and healthcare administrators in enhancing and enforcing programs that train doctors on resilience skills.

INTRODUCTION

In recent decades, burnout has become a global epidemic affecting various professions. The burnout is defined by Maslach et al. (2001) as a reaction to sustained work-related stress that depends on the level of emotional depletion, cynicism, and professional efficacy. It primarily affects work accomplishment, job satisfaction, interpersonal relations, and makes one more vulnerable to illness (Chemali et al., 2019). Resilience, which is the capacity to endure stress (Connor and Davidson, 2003), is a protective measure against burnout and is highly associated with life satisfaction (Aboalshamat et al., 2018), helping in maintaining greater well-being.

Doctors have a crucial part in patient care ensuring the proper health of populations. But in many cases, the doctors are exposed to extreme levels of occupational stressors and burnouts due to the nature of the work they undergo. These harsh factors can catastrophically impact their quality of life (Wu et al., 2007). A study carried out on the doctors nationwide in the US over a six-month period revealed that 44.0% of these

DOI: <http://doi.org/10.5281/zenodo.20569901>

doctors had one or more symptoms of burnout in their career (Harry et al., 2020). Another study by the European General Practice Research Network in 2008 reported that burnout was also a recurrent issue across Europe (Soler et al., 2008). Several studies showed that burnout was highly prevalent among doctors and healthcare professionals across the world, including China, Poland, and Italy (Lo et al., 2018; Misiotek et al., 2015; Sanfilippo et al., 2018).

According to the recent survey (American Medical Association & American Medical Association, 2025), the prevalence of burnout is highest among the departments: Emergency medicine (52.2%), Family medicine (46.4%), Obstetrics and gynecology (45.8%), Pediatrics (42.1%), Internal medicine (42%), Hospital medicine (40.6%). In another study, when compared with doctors in other departments (38%) and the general population (30%), Emergency Medicine doctors were particularly vulnerable to burnout (Arora et al., 2013). In the same way, the burnout rates are so much higher in orthopedic surgery because its training is extremely challenging, because of the necessity both to increase knowledge and to learn surgical techniques (Daniels et al., 2016).

Particularly, the doctors in Pakistan are more susceptible to burnout. The rates of burnout and stress are highly prevalent among house officers, residents, surgeons, and nurses due to low salaries, high work demands, and a lack of support from colleagues (Liaqat et al., 2019; Mazhar et al., 2019). Many studies report that numerous factors are coupled with burnout, including time strain, role conflicts, lack of support from work-fellows, high patient-to-staff ratios, frequent contact with terminally sick patients, long operating hours, and frequency of on-call duties (Ahmed & Noushad, 2013). Thus, doctors experience psychosocial stressors that create a stressful environment, eventually resulting in burnout

DOI: <http://doi.org/10.5281/zenodo.20569901>

syndrome and other maladaptive behaviors that impact their well-being and reduce productivity (Gómez et al., 2019).

According to the recent studies, psychological resources are important in determining how an individual is perceived to control the challenging and stressful environment (Lenzenweger, 2004). The personality factor that has been studied as a key predictor of burnout is resilience (García and Calvo, 2011). Resilience is a buffering mechanism against the adverse impact regarding stress and reduces burnout (Guo et al., 2017). A study explains that resilience is one element that may help to explain why certain doctors can manage stress well and recover from stressful situations (West et al., 2020). According to West et al. (2020), a large-scale study on 5445 US doctors reported that resilience is negatively associated with symptoms of exhaustion. Highly resilient doctors are more expected to adapt to tough situations and pressurized tasks in clinical settings (Epstein & Krasner, 2013). Several studies have also suggested that resilience might safeguard individuals from severe stress, and it inversely impacts the burnout and depressive symptoms among hospital personnel (Di Giuseppe et al., 2021; Meynaar et al., 2020).

Researches in resident doctors demonstrate that those doctors who are more resilient reported less burnout in both medical and surgical specialties. In a relevant example, a U.S. national survey of residents reported a positive relationship between resilience, program support, and autonomy, and a negative relationship between resilience and burnout in both medical and surgical trainee groups (Nituica et al., 2020). Similar strong negative associations have been found between resilience (and grit) and burnout among surgeons in surgical specialties like orthopedics (Hamdan et al., 2023). Therefore, developing resilience among doctors should be prioritized.

DOI: <http://doi.org/10.5281/zenodo.20569901>

According to Shi et al. (2015), life satisfaction, positive affect, and negative affect make up positive well-being. Similarly, life satisfaction is a significant factor of positive well-being (Vittersø, 2025), and it measures how individuals judge their lives against self-identified criteria and against what they possess. Health-care worker satisfaction, in particular, affects the productivity, quality, efficiency, and dedication to work and, at the same time, health care costs. (Koftuniuk et al., 2021). Stress and burnout have a detrimental effect on life quality and can result in a number of unfavorable consequences. According to some studies, younger doctors who perform high-risk procedures and experience work-life conflicts are at greatest risk (Lacy & Chan, 2017). Factors that are frequently related to higher burnout include younger age, being female, overloaded work-related tasks, long working hours, specialty types (e.g., emergency medicine, high-acuity fields), and insufficient support (Ashraf et al., 2020). Factors that frequently influence life satisfaction among doctors are daily work and home stressors, job requirements, and peer relationships. Doctors who tend to have a good work-life balance and better support from coworkers report higher life satisfaction (Mahmood et al., 2019).

LITERATURE REVIEW

Burnout is an “occupational phenomenon”, not a medical condition, that consists of emotional exhaustion, physical tiredness, and cognitive fatigue (Maslach & Leiter, 2001). Constant burnout is the reason behind diminished quality of life and is linked to increased vulnerability to sleep-related issues as well as to numerous medical conditions such as mild cognitive impairment, diabetes, and cardiovascular disease (Grossi et al., 2015). Burnout has become a global crisis, especially among doctors, it has reached epidemic levels. The healthcare industry is posing several stressors on healthcare professionals, just

DOI: <http://doi.org/10.5281/zenodo.20569901>

like other professions, including numerous problems from meeting time constraints to unmanageable work procedures, incompatibility with seniors, challenging demands, and the burdens of clinical work (Fitzpatrick et al., 2019). The syndrome of professional burnout has been reported among physicians in various countries (Romani & Ashkar, 2014) and in most fields of medical practice, such as general medicine, surgery, intensive care, surgeons, oncologists, and anesthetists (Shanafelt, 2009)

The concept of burnout was presented first in the occupational field. It was an American Psychologist Herbert Freudenberger who coined this term and he described it as being the extinction of motivation or incentive (Freudenberger, 1974). In the conceptualization of burnout, Maslach and Jackson defined it as a three-dimensional construct which incorporated the emotional exhaustion, depersonalization, and lack of personal accomplishment (Maslach et al., 2001). Burnout has been formally recognized by the WHO as a syndrome caused by chronic stress related to the workplace that has not been effectively managed and was included in the 11th Revision of the International Classification of Diseases (World Health Organization: WHO, 2019).

Burnout is also well-known issue that affects work performance, job satisfaction, relationships with other people, and susceptibility to illnesses (Chemali et al., 2019). Burnout has also been associated with anxiety symptoms, low job satisfaction, and a higher risk of medical errors (Lo et al., 2018). It also plays a major role in deteriorating health, resulting in headaches, sleep issues, exhaustion, marital and familial issues, anxiety, depression, high blood pressure, and may lead to alcoholism and drug abuse (Maslach et al., 2001). Moreover, Irregular shift work, including working hours and night work, which disturbs the circadian rhythm, also has an adverse effect (Basile, 2021).

DOI: <http://doi.org/10.5281/zenodo.20569901>

Burnout, a global issue, has long-term effects on doctors' careers, patient care and safety, medical workforce stability, and the healthcare system (West et al., 2006; Shanafelt et al., 2009; Liu et al., 2024). The risk of patient safety incidents has doubled in case of doctor burnout (Hodkinson et al., 2022), and the latter is strongly correlated with such negative outcomes as medical errors, increased costs incurred by health care professionals, and negative health outcomes and poor quality of care in the long run (Moss et al., 2016; Pastores et al., 2019; Sinbukhova et al., 2019). According to the literature, Lacy and Chan (2017) report that increased burnout levels are associated with early retirement, increased job change, and clash of marital or relationship with increased depression, alcohol abuse, drug abuse, and suicide. Physician burnout, therefore, is a significant issue in medical and even in the health context.

Resilience is an important psychological resource that has attracted a lot of attention in the medical field by the researchers. The general definition of resilience is the capacity to recover or survive in a situation and therefore record beneficial results even after an unpleasant circumstance or experience (Vella & Pai, 2019). According to Connor and Davidson, resilience is a personal attribute that determines individuals to succeed in the face of adversities (Connor and Davidson, 2003). In the framework of resilience among doctors, it is viewed as a dynamic process and prompt response to various challenges (Sheikhrabori et al., 2022). The resilient doctors do not mean that they do not feel or experience grief, anxiety, or stress; it is only that they apply their best coping strategies to cope with the crisis and live to cope with the challenges (Cherry, 2022).

In psychology, the concept of resilience emerged in the field of developmental psychology. Emmy Werner, developmental psychologist, in her popular research of children being raised in stressful conditions discovered that some children managed to

DOI: <http://doi.org/10.5281/zenodo.20569901>

succeed even despite the difficulties encountered (E. Werner, 1997). These children also exhibited what Werner referred to as resilient behavior, they adjusted themselves positively to adversity and learned to cope. The American psychological association (2014) defines resilience as a process of adapting effectively to misfortune, trauma, tragedy, danger or even a major stressful experience.

Highly resilient individuals were observed as persevering, self-reliant, and equanimous, and they possessed explicit goals in life which they were working tirelessly to attain despite frustrations and obstacles (G. M. Wagnild and Collins, 2009). It assists individuals in acquiring coping mechanisms, reduces anxiety, and gain skills in problem-solving (Rushton et al., 2015). According to Ferreira and Gomes (2021), resilience plays a key role in buffering against the intense stressors, resulting in decreased levels of burnout and increased personal accomplishment. Thus, resilience acts as a protective factor. Resilience results in enhanced patient care quality, professional autonomy, and increased job satisfaction (Fazekas et al., 2024).

Resilience not only enables doctors to combat stress but also to bounce back from it (Aburn et al., 2016). Resilience in primary care settings was identified to be a combination of personal aspects with social and workplace characteristics (Robertson et al., 2016). Personal resilience helps doctors to adapt to sudden changes in patient condition, emergencies, and high-stakes choices within the medical profession, thereby facilitating the preservation of safe patient care (W. Wang et al., 2024). For doctors, whose duties often involve life-threatening consequences, ethical stress, and fast responses, resilience improves the ability to cope, educate, ponder, and recover, which subsequently reduces the burnout risk and enhances well-being (Haraldseid-Driftland et al., 2022). Highly resilient people have enhanced coping to work stress and difficulties, hence enhancing

DOI: <http://doi.org/10.5281/zenodo.20569901>

performance (Abro et al., 2023). Thus, doctors' resilience reduces burnout and enhances mental health.

Life satisfaction has been described as the cognitive and affective assessments of an individual concerning his or her life (Diener et al., 2009). Life satisfaction is an important determinant of the overall well-being and, therefore, a key predictor of psychological well-being (Lombardo et al., 2018). Wijayatunga (2021) found that the levels of life satisfaction were much higher among healthcare professionals with higher status, postgraduate qualifications, higher income, and perceived social support. An additional cross-sectional survey of nurses in Türkiye further revealed that positive working conditions were linked with positive satisfaction with life (Medeni et al., 2025).

In the 1980s, Ed Diener suggested a systematic model of subjective well-being that consists of three primary components, namely, (1) positive affect, (2) negative affect, and (3) life satisfaction. Despite the fact that the two former are emotion-related, the third, life satisfaction, is the cognitive aspect of subjective well-being (MSEd, 2023). According to Diener et al. (2009), the level of positive evaluation of the quality of life by a person is defined as life satisfaction. Positive Psychology, led by researchers like Martin Seligman, has changed life satisfaction to be treated alongside flourishing, meaning, and purpose, rather than just hedonic happiness alone (Petrakova, n.d.). The existing empirical data reveal that high life satisfaction correlates with better health outcomes, less stress, better coping skills, fewer visits to the doctor, and numerous other positive implications (Kim et al., 2013). However, doctors often face long working hours, rotating shifts, high stress, professional ethics, and poor working conditions, which all impact life satisfaction (Mahmood et al., 2019).

DOI: <http://doi.org/10.5281/zenodo.20569901>

Many studies report that individuals with higher life satisfaction tend to show positive work-related behaviors, i.e, helping coworkers, going beyond formal job duties, lower emotional exhaustion, and more trust in colleagues (Merkin, 2023). Therefore, it describes that higher life satisfaction is also linked with better performance, commitment, and lower turnover intention for organizations. Organizations benefit when employees express satisfaction with their life and work-life balance through reduced absenteeism, increased retention, and engagement, which is the foundation of productivity and the long-term sustainability of organizations (Leitão et al., 2019). Thus, when life satisfaction is higher, doctors are better able to provide quality patient care, maintain professional standards. Burnout has captured a great part of scholarly attention. Many studies have tended to show a significant negative relationship between burnout and life satisfaction (Łaskawiec-żuławińska et al., 2024; Shim & Go, 2025). For example, in a recent study on medical students (Wang et al., 2022), burnout had negatively predicted life satisfaction. Additionally, the association between burnout and life satisfaction among doctors has not been extensively studied. Many empirical findings reveal an inverse relationship between resilience and burnout. For example, Alonazi et al. (2023) found a negative correlation between psychological resilience and burnout among nurses. Moreover, McCain et al. (2017) report that higher burnout was linked with lower resilience among doctors in the UK, leading to higher use of maladaptive coping skills. A systematic review of surgeons also identified resilience as a buffer against the negative impacts of burnout and work-related stressors (Otukoya et al., 2025).

Extensive empirical data indicate a positive association between resilience and life satisfaction, where higher resilience tends to predict higher life satisfaction. Greater population studies (Karataş & Tagay, 2020) and longitudinal studies (Wang et al., 2022)

DOI: <http://doi.org/10.5281/zenodo.20569901>

confirm that resilience is positively correlated with life satisfaction. Secondary outcomes or related well-being constructs, such as quality of life, sense of well-being, and professional quality of life, are also often positively associated with resilience (Fazekas et al., 2024). For instance, doctors who were more resilient experienced fewer burnout symptoms and improved satisfaction with or perceived quality outcomes (McCain et al., 2017). Hence, resilience is generally seen as a factor that promotes subjective well-being (including life satisfaction). Despite an expanding body of research on doctors' well-being, including the variables resilience, burnout, and life satisfaction, several significant gaps remain, especially in the context of doctors working with high-risk patients in the Medicine and Surgery departments.

Few studies have simultaneously measured resilience, burnout, and life satisfaction within the same model and much of the literature is concentrated among nursing, medical students, or healthcare professionals, especially in high-income countries. Systematic reviews show that the majority of burnout/stress research focuses on nurses rather than physicians (Lozano et al., 2021). Moreover, none of the studies measure these relationships among various high-risk specialties, like Medicine versus Surgery. Although most studies report burnout in medicine, emergency, surgery, and psychiatry separately (Anjum et al., 2023; McCain et al., 2017; Zhang et al., 2020; Murtaza, 2025) but few studies compare Medicine and Surgery departments in the same sample. Lastly, there is insufficient measurement of global life satisfaction among doctors. Although most of the available literature focuses on quality of life, job satisfaction, or work-life balance, a smaller number of studies specifically quantify life satisfaction with validated measures (Sarwar et al., 2023).

DOI: <http://doi.org/10.5281/zenodo.20569901>

These gaps are more intense in the Pakistani context. Although research has documented high burnout among doctors in various specialties in Pakistan (Hussain et al., 2022), there are very limited studies that include resilience as a protective factor and life satisfaction as an outcome in high-risk specialties such as Surgery and Medicine. Comparative studies are lacking between these departments, and resilience and life satisfaction metrics are barely validated in Pakistani doctors. In order to fill these gaps, the proposed research will investigate the relationships between resilience, burnout, and life satisfaction among the doctors involved in dealing with high-risk patients in the Medicine and Surgery departments. The study will also offer new insights on the connection between resilience and life satisfaction with burnout in high-risk clinical settings in Pakistan by measuring all three variables at the same time and comparing them across these two high-need specialties.

Hypotheses of the study

1. There will be a significant positive correlation between resilience and life satisfaction.
2. There will be a significant negative correlation between burnout and life satisfaction.
3. There will be a significant negative correlation between resilience and burnout.
4. Resilience and burnout will jointly and significantly predict life satisfaction among doctors dealing with high-risk patients.
5. Female doctors will have higher burnout levels than male doctors.
6. Married doctors will report higher resilience than those who are unmarried.

DOI: <http://doi.org/10.5281/zenodo.20569901>

7. Doctors in the Surgery department report a higher level of burnout than those in the Medicine department.

METHODOLOGY

Sample

A quantitative, cross-sectional, correlational research was conducted in the surgery and medical department of Allied 1 and 2 Hospital, Faisalabad. It involved the doctors dealing with high-risk patients. The study sample consisted of 151 medical doctors involved in the care of high-risk patients (77 female, 74 male), ranging in age from 22 to 32 ($M = 26.32$, $SD = 2.86$). Sixty percent were unmarried. The breakdown of medical specialty was: 45.7% for surgery and 54.3% for medicine. Sixty-two percent were house-officers, and thirty-seven percent were residents.

Demographics

Demographic variables such as age, gender, marital status, monthly income, current position, years of experience and on-call frequency were collected.

Copenhagen Burnout Inventory (CBI)

This 19-item scale, developed by Kristensen et al., measured the burnout among doctors. It assessed burnout in three different aspects: personal burnout, work related burnout, and patient related burnout. All items are scored on a 5-point Likert scale (Always, Often, Sometimes, Seldom, Never) and computed to average the responses on each scale, where higher scores indicate higher burnout. Internal reliability in the present study was $\alpha = 0.79$ for the Overall Copenhagen Burnout Inventory, $\alpha = 0.66$ for Personal Burnout, $\alpha = 0.67$

DOI: <http://doi.org/10.5281/zenodo.20569901>

for work-related burnout (without recoding the items), and $\alpha = 0.74$ for patient-related Burnout.

Brief resilience scale (BRS)

The 6-item Brief Resilience Scale was used to measure resilience (Smith et al., 2008). It assesses how people bounce back to life after stress. The items are evaluated on a 5-item Likert scale (where 1 = strongly disagree, 5 = strongly agree) and will be summed up (with a recoding of three items) to give scores with higher scores indicating greater resilience. In the current study, internal reliability was = 0.50.

Satisfaction with Life Scale

This five-item Satisfaction with Life Scale (SWLS) scale is based on the subjective assessment of the sense of life satisfaction developed by Diener et al. It measured global cognitive judgments of one's life satisfaction on a 7-point Likert scale from 1 (strongly disagree) to 7 (strongly agree). The Cronbach's alpha value for this scale is 0.80.

High-risk Patients

In the conceptualization of high-risk patients, doctors from the medicine and general surgery departments of the hospital were included, as these specialties routinely care for patients with critical, life-threatening conditions. "High-risk patients" are defined as those with severe, acute, or complex medical problems requiring intensive monitoring or urgent intervention. For example, in the field of surgery, emergency patients are specifically defined as the high-risk group because of the elevated morbidity and mortality (Phillips et al., 2019). By extension, we consider any patient needing intensive or emergency care,

DOI: <http://doi.org/10.5281/zenodo.20569901>

such as multi-system organ failure, acute trauma, or post-operative critical care, to be “high-risk” in our study context (Phillips et al., 2019).

Doctors from medicine and general surgery were included specifically because these departments regularly treat high-risk patients. In other words, doctors in these specialties face similar stressors to those documented in “high-stress” units (Filaj et al., 2025). Medicine wards (e.g. managing sepsis, cardiac crises) and surgical wards (e.g. managing perioperative critical cases) involve analogous high-acuity cases, justifying this study focus on these departments. Likewise, caring for critically ill patients is described as a highly emotionally charged work environment; doctors in such settings are known to be at elevated risk of burnout (Ciuffini et al., 2022).

Procedure

This current study was conducted in the medicine and surgery departments of Allied 1 and 2 hospitals, Faisalabad. The doctors working in these departments were invited to participate in the research. The researcher in-person visited each department and approached those doctors fulfilling the inclusion criteria. They were informed about the study objectives, confidentiality, and voluntary participation. They were also informed of their right to withdraw at any time without consequences. Those who agreed were given questionnaires to complete on the spot, while busy doctors were given a copy of the questionnaire to return within the time frame of 3 days. Written Informed Consent was also obtained from each participant

DOI: <http://doi.org/10.5281/zenodo.20569901>

RESULTS

Table 1

Descriptive Statistics and Alpha Reliabilities of All Study Variables (N= 151)

Scales	<i>k</i>	<i>M</i>	<i>SD</i>	<i>a</i>	Potential Range	Actual Range	Skewness
BRS	6	17.24	3.29	0.50	6-30	6-25	-0.19
SWLS	5	20.96	6.40	0.80	5-35	7-35	0.09
CBI	19	52.30	13.37	0.79	0-100	18-89	0.18
Personal Burnout	6	55.71	16.13	0.66	0-100	8-91	0.13
Work-related Burnout	7	52.50	13.24	0.67	0-100	8-85	-0.29
Patient-related Burnout	6	48.60	19.72	0.74	0-100	0-100	0.02

Note. BRS = brief resilience scale; SWLS = satisfaction with life scale; CBI = Copenhagen burnout inventory

Table 1 postulates mean, standard deviation, alpha reliability, and skewness. Values of all variables examined in the current study.

DOI: <http://doi.org/10.5281/zenodo.20569901>

Table 2

Correlational Matrix for All the Variables Used in The Study (N=151)

Variable	1	2	3	4	5	6
1. BRS	–					
2. SWLS	0.42***	–				
3. CBI	-0.18*	-0.36***	–			
4. PB	-0.10	-0.35***	0.78***	–		
5. WB	-0.12	-0.26**	0.83***	0.46***	–	
6. CB	-0.20*	-0.28***	0.87***	0.48***	0.59***	–

Note. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, BRS= Brief Resilience Scale, SWLS= Satisfaction with Life Scale, CBI= Copenhagen Burnout Inventory, PB= Personal Burnout, WB= Work-related Burnout, CB= Patient-related Burnout

Table 2 explains the intercorrelations among scales and subscales. The results shows Pearson correlation between resilience (BRS), burnout (CBI) and life satisfaction (SWLS).

Table 3

Hierarchical Multiple Regression predicting life satisfaction from Resilience and Burnout (N=151)

Variables	B	SE	β	t	R ²	ΔR^2
Step 1						
Resilience	0.81	0.15	0.42	5.59***	0.17	0.16
Step 2						
Resilience	0.75	0.14	0.39	5.47***	0.28	0.27

DOI: <http://doi.org/10.5281/zenodo.20569901>

Personal	-0.02	0.01	-0.32	-4.56***
Burnout				

Note Step 1: $F(1, 151) = 31.28, p < 0.001$. Step 2: $F(2, 151) = 28.14, p < 0.001$

Table 3 shows Multiple Linear Regression indicating Resilience, Burnout and its subscales as predictors of life satisfaction among doctors dealing with high-risk patients.

Table 4

T- Test analyses were carried out to seek the gender differentiation (i.e., males, females) with reference to study variables.

Scales & subscales	Males (n = 74)		Females (n = 77)		t(151)	95% CI		Cohen's d
	M	SD	M	SD		LL	UL	
BRS	16.71	3.21	17.74	3.30	-1.93	-2.07	0.02	0.31
SWLS	21.11	7.12	20.79	5.66	0.30	-1.76	2.39	0.05
CBI	50.49	14.28	53.96	12.29	-1.60	-7.76	0.82	0.26
PB	54.05	17.04	57.31	15.13	-1.24	-8.44	1.94	0.20
WB	50.29	14.39	54.59	11.75	-2.01*	-8.54	-0.06	0.33
CB	47.18	20.83	49.95	18.63	-0.86	-9.13	3.60	0.14

Note. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, BRS= Brief Resilience Scale, SWLS= Satisfaction with Life Scale, CBI= Copenhagen Burnout Inventory, PB= Personal Burnout, WB= Work-related Burnout, CB= Patient-related Burnout

DOI: <http://doi.org/10.5281/zenodo.20569901>

The above table indicates the mean, standard deviation, and t-test value for males and females on the Brief Resilience Scale, Satisfaction with Life Scale, Copenhagen Burnout Inventory, and its subscales.

Table 5

T- Test analyses were carried out to seek the marital status differentiation (i.e., married, unmarried) with reference to study variables

Scales & subscales	Married (n = 57)		Unmarried (n = 91)		t(151)	95% CI		Cohen's d
	M	SD	M	SD		LL	UL	
BRS	16.88	3.61	17.94	2.60	-2.09*	-2.08	-0.06	0.33
SWLS	21.81	5.99	20.60	6.64	-1.14	-3.29	0.88	0.19
CBI	52.43	10.31	51.76	14.84	-0.32	-4.76	3.42	0.05
PB	55.41	11.27	55.67	18.65	0.11	-4.59	5.13	0.02
WB	52.81	11.92	52.05	14.06	-0.35	-5.03	3.51	0.06
CB	49.04	16.48	47.57	21.13	-0.47	-7.62	4.67	0.07

Note. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, BRS= Brief Resilience Scale, SWLS= Satisfaction with Life Scale, CBI= Copenhagen Burnout Inventory, PB= Personal Burnout, WB= Work-related Burnout, CB= Patient-related Burnout

The above table indicates the mean, standard deviation, and t-test value for marital status differentiation on the Brief Resilience Scale, Satisfaction with Life Scale, Copenhagen Burnout Inventory, and its subscales

Table 6

T- Test analyses were carried out to seek the department-wise differentiation (i.e., medicine, surgery) with reference to study variables

Scales & subscales	Medicine ($n = 82$)		Surgery ($n = 69$)		$t(151)$	95% CI		Cohen's d
	M	SD	M	SD		LL	UL	
BRS	17.51	3.51	16.91	2.99	1.13	-0.44	1.64	0.18
SWLS	20.64	6.48	21.30	6.32	-0.63	-2.72	1.41	0.10
CBI	53.72	12.66	50.52	14.07	1.46	-1.14	7.54	0.24
PB	58.74	15.63	52.11	16.07	2.55*	1.50	11.75	0.42
WB	53.44	12.56	51.33	14.01	0.96	-2.20	6.43	0.16
CB	49.09	19.88	48.01	19.66	0.33	-5.30	7.46	0.05

Note. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, BRS= Brief Resilience Scale, SWLS= Satisfaction with Life Scale, CBI= Copenhagen Burnout Inventory, PB= Personal Burnout, WB= Work-related Burnout, CB= Patient-related Burnout

The above table indicates the mean, standard deviation, and t-test value for medicine and surgery departments on the Brief Resilience Scale, Satisfaction with Life Scale, Copenhagen Burnout Inventory, and its subscales.

DISCUSSION

This study focuses on investigating the association between burnout, resilience, and life satisfaction among doctors dealing with patients at high-risk in Medicine and Surgery Departments. The results largely supported the hypothesized relationships.

The results supported hypothesis 1, showing a statistically significant positive relationship between resilience and life satisfaction, which aligns with the growing body of research (N. Wu et al., 2022; Karagöz et al., 2021; Khalid, 2021; Sclanavo, 2023; Aboalshamat et al., 2018). In line with this study, West et al. (2020) showed that doctors facing high-risk patient care demands, it appears that those who are more resilient are better able to maintain a satisfying life, and perhaps buffer the impact of work stress on their personal life. The findings align with the Conservation of Resources (COR) theory (Hobfoll, 2010). According to this theory, resilience acts to safeguard against resource loss, promoting positive outcomes and therefore enhancing life-being.

The results of the current study confirm the hypothesis 2 that resilience and burnout are negatively related to each other among health care providers who work with high-risk patients, which proves empirical evidence in healthcare environments (Castillo-González et al., 2023; Zahednezhad et al., 2021; J. L. Jackson et al., 2023; Amiri et al., 2024). It states that the greater the resilience of doctors working with high-risk patients, the lower the burnout. Stress, according to the COR theory (Hobfoll, 2010), is caused by the threat to

valuable personal resources. Nevertheless, resilient doctors can better withstand their resources and therefore defend themselves against resource breach.

This finding supported hypothesis 3, indicating that burnout has a statistically significant negative correlation with life satisfaction, which corresponds well with the emerging body of research, especially in healthcare and other high-stress groups (Łaskawiec-Żuawińska et al., 2024; Marzo et al., 2022; Uchmanowicz et al., 2019). This demonstrates that physicians with high burnout rates when handling high-risk patients have higher chances of reporting lesser levels of life satisfaction. In reference to COR theory (Hobfoll et al., 2010), in case of constant stressors, people experience widespread loss of resources. It usually leads to adverse consequences, i.e., burnout, which in its turn causes reduced life satisfaction.

This study also analyzed the effect of resilience and burnout on life satisfaction. The multiple regression model was used, and the results found that both variables had a significant predictive effect on life satisfaction, explaining 28% variance in life satisfaction. In this model, resilience was a significant positive predictor of life satisfaction, suggesting that resilience acts as a protective factor for doctors working with high-risk patients and therefore is a key determinant of well-being. This finding is consistent with existing literature (Gonzalez et al., 2025; Sultana et al., 2024). This can be explained by the fact that resilience provides people with the capacity to persevere through hardship (Sclanavo, 2023) and acts to safeguard against life challenges and difficulties (M. Liu et al., 2024).

In contrast, burnout was found to be a significant negative predictor of life satisfaction ($R^2 = -0.28$, $p < 0.001$), suggesting that higher burnout level has a significant effect on lower life satisfaction. These results supports the existing literature (Q. Wang et al., 2022; Permarupan et al., 2020; Shim & Go, 2025). While burnout does affect life satisfaction, not

all burnout domains contributed. This study's results showed that only personal burnout was a significant predictor of life satisfaction, rather than work or patient burnout. It suggests that the internal, individual experience of burnout may matter most for life satisfaction in this context of high-risk demands. Therefore, the internalized experience of burnout has the greatest impact on overall life satisfaction (De Diego-Cordero et al., 2021). It was hypothesized in hypothesis 5 that female doctors would report higher overall burnout than the males. While the results showed that significant gender difference emerged on the work-related burnout subscale. This indicates that female doctors in our sample experience a greater burden in the work domain. This could be explained in light of gender-specific stressors such that female doctors are exposed to greater job demands (Lyubarova et al., 2023) leading to an increase in work-related burnout. Moreover, dual responsibilities, including professional duties along with domestic/caregiver responsibilities (Marshall et al., 2020), and gender-related discrimination (Rotenstein et al., 2021) suggest a reasonable explanation.

The results did not support hypothesis 6 that married doctors tend to be more resilient than the unmarried. Only one prior study, conducted on students, similarly reported lower resilience among married individuals, indicating limited supporting literature. (Nijimbere et al., 2017). These results may be interpreted as married doctors have more family and marriage-related burdens, including child-care, family and household management (Zatuski & Makara-Studzińska, 2024). Unmarried doctors, on the contrary, might have more flexibility and autonomy, which allows them to concentrate on the development of coping strategies and self-management (Mansi et al., 2025). Moreover, in the context of Pakistan, the marital role is also characterized by additional non-work demands that can further worsen the work-family conflict and diminish resilience (Misfin et al., 2024).

The study results did not support hypothesis 7 that doctors working in the surgery department will exhibit higher burnout than those working in the medicine department. Instead, the data revealed that doctors working in the medical department reported a higher level of burnout. Low et al. (2019) in a meta-analysis found that burnout among medical and surgical residents is higher, but the difference is not significant. Therefore, the results of this study suggest that many factors can be found to play a contributing role such that doctors in the medical department often deal with chronically ill at-high-risk patients, with slower progress, and have a high cognitive burden (Iskander, 2018). Moreover, cultural factors in Pakistani doctors also play a key role, such that the hierarchical norms, high burden of patients, and limited institutional support have been identified as contributors to doctor burnout in Pakistan (Hashmi et al., 2021). Therefore, all these factors contribute to emotional strain and burnout in doctors and appear to be the reasons behind a higher level of burnout in the Medicine department.

Limitations and Recommendations

The sample used in this research is limited to doctors who treated high-risk patients in the Medicine and Surgery Department Allied 1 and 2 Hospitals at Faisalabad. The duration was restricted and informal conclusions are not guaranteed since it was a correlational study. The data collection was done through self-report questionnaires only and this can lead to common method variance, social desirability or response bias.

To be more generalizable, further studies might increase sampling to other specialties, hospitals, and regions. Mediation or moderation analysis can also be incorporated into future studies. A qualitative study is needed to understand the impact of resilience in

doctors on burnout and its impact on their life satisfaction. Longitudinal study can also be applied in establishing the causal path or bidirectional relationship.

Conclusion

The purpose of this study was to explore the association between resilience, burnout, and life satisfaction among doctors dealing with high-risk patients in the department of medicine and surgery. As per the findings of this study, resilience and personal burnout are key determinants of life satisfaction among doctors. Resilience acts as a protective factor, while personal burnout acts as a risk factor. Although other burnout dimensions showed correlation with life satisfaction, their predictive utility was not robust when controlling for personal burnout and resilience. There were also group differences in demographic comparisons, such as that females rated higher in work-related burnout, unmarried doctors rated higher in resilience, and medicine doctors rated higher in personal burnout than surgeons. These observations help to isolate those areas of burnout that would be most pertinent to well-being in the medical profession. The provided results also indicate the necessity to reinforce the resilience and address personal burnout in an effort to facilitate life satisfaction among medical workers and physicians in high-stress environments.

REFERENCES

- Aboalshamat, K., Alsiyud, A., Al-Sayed, R., Alreddadi, R., Faqiehi, S., & Almehmadi, S. (2018). The relationship between resilience, happiness, and life satisfaction in dental and medical students in Jeddah, Saudi Arabia. *Nigerian Journal of Clinical Practice*, 21(8), 1038. https://doi.org/10.4103/njcp.njcp_278_17

- Abro, S., Chanar, Z., Sikandar, P., & Rahim, S. (2023). The Power of commitment: How can high commitment work system build resilience, combat stress and cynicism. *Pakistan Journal of Commerce and Social Sciences*, 17(1), 39. <https://doi.org/10.64534/commer.2023.086>
- Aburn, G., Gott, M., & Hoare, K. (2016). What is resilience? An Integrative Review of the empirical literature. *Journal of Advanced Nursing*, 72(5), 980–1000. <https://doi.org/10.1111/jan.12888>
- Ahmed, S., & Noushad, S. (2013). Sorts and sources of stress in Pakistan; A Comprehensive outlook. *INTERNATIONAL JOURNAL OF ENDORSING HEALTH SCIENCE RESEARCH (IJEHSR)*, 1(1), 04. <https://doi.org/10.29052/ijehsr.v1.i1.2013.4-8>
- Alonazi, O., Alshowkan, A., & Shdaifat, E. (2023). The relationship between psychological resilience and professional quality of life among mental health nurses: a cross-sectional study. *BMC Nursing*, 22(1). <https://doi.org/10.1186/s12912-023-01346-1>
- American Medical Association & American Medical Association. (2025, May 13). Which physician specialties are seeing a drop in burnout? *American Medical Association*. <https://www.ama-assn.org/practice-management/physician-health/which-physician-specialties-are-seeing-drop-burnout>
- American Psychological Association. (2014). The road to resilience. Washington, DC: American Psychological Association. Retrieved from <http://www.apa.org/helpcenter/road-resilience.aspx>
- Amiri, S., Mahmood, N., Mustafa, H., Javaid, S. F., & Khan, M. A. (2024). Occupational Risk Factors for Burnout Syndrome Among Healthcare Professionals: A Global Systematic Review and Meta-Analysis. *International Journal of Environmental Research and Public Health*, 21(12), 1583. <https://doi.org/10.3390/ijerph21121583>

- Anjum W, Sajid A, Rafiq M, Chaudhary NI, Rabab E (2023). Burnout, Spiritual Well-Being, And Life Satisfaction Among General Surgeons of Tertiary Care Hospital Lahore. *History of Medicine* 9(2): 705–711. <https://doi.org/10.17720/2409-5834.v9.2.2023.089>
- Arora, M., Asha, S., Chinnappa, J., & Diwan, A. D. (2013). Review article: Burnout in emergency medicine physicians. *Emergency Medicine Australasia*, 25(6), 491–495. <https://doi.org/10.1111/1742-6723.12135>
- Ashraf, F., Ahmad, H., & Aftab, S. (2020). Interactive role of personal and work-related factors in psychological burnout: a study of Pakistani doctors. *Journal of the Pakistan Medical Association*, 0, 1. <https://doi.org/10.5455/jpma.35303>
- Basile, M. (2021). Prevalence of stress, burnout syndrome, anxiety and depression among physicians of a teaching hospital during the COVID-19 pandemic. *Archivos Argentinos De Pediatría*, 119(5). <https://doi.org/10.5546/aap.2021.eng.317>
- Castillo-González, A., Velando-Soriano, A., De La Fuente-Solana, E. I., Martos-Cabrera, B. M., Membrive-Jiménez, M. J., Lucía, R., & La Fuente, G. a. C. (2023). Relation and effect of resilience on burnout in nurses: A literature review and meta-analysis. *International Nursing Review*, 71(1), 160–167. <https://doi.org/10.1111/inr.12838>
- Chemali, Z., Ezzeddine, F. L., Gelaye, B., Dossett, M. L., Salameh, J., Bizri, M., Dubale, B., & Fricchione, G. (2019). Burnout among healthcare providers in the complex environment of the Middle East: a systematic review. *BMC Public Health*, 19(1). <https://doi.org/10.1186/s12889-019-7713-1>
- Cherry, K. (2022). What is Resilience. Verwellmind. <https://www.verywellmind.com/what-is-resilience-2795059>

- Ciuffini, R., Marrelli, A., Leuter, C., Stratta, P., Paladini, A., Ciccozzi, A., Marsili, I., Marinangeli, F., & Piroli, A. (2022). The stress response of intensive care unit medical doctors facing repeated severe emergencies. *Frontiers in Psychology, 13*, 895954. <https://doi.org/10.3389/fpsyg.2022.895954>
- Connor, K. M., & Davidson, J. R. (2003). Development of a new resilience scale: The Connor-Davidson Resilience Scale (CD-RISC). *Depression and Anxiety, 18*(2), 76–82. <https://doi.org/10.1002/da.10113>
- Daniels, A. H., DePasse, J. M., & Kamal, R. N. (2016). Orthopaedic surgeon burnout. *Journal of the American Academy of Orthopaedic Surgeons, 24*(4), 213–219. <https://doi.org/10.5435/jaaos-d-15-00148>
- De Diego-Cordero, R., Iglesias-Romo, M., Badanta, B., Lucchetti, G., & Vega-Escañó, J. (2021). Burnout and spirituality among nurses: A scoping review. *EXPLORE, 18*(5), 612–620. <https://doi.org/10.1016/j.explore.2021.08.001>
- Di Giuseppe, M., Nepa, G., Prout, T. A., Albertini, F., Marcelli, S., Orrù, G., & Conversano, C. (2021). Stress, Burnout, and Resilience among Healthcare Workers during the COVID-19 Emergency: The Role of Defense Mechanisms. *International Journal of Environmental Research and Public Health, 18*(10), 5258. <https://doi.org/10.3390/ijerph18105258>
- Diener, E., Emmons, R. A., Larsen, R. J., & Griffin, S. (1985). The satisfaction with life scale. *Journal of Personality Assessment, 49*(1), 71–75. https://doi.org/10.1207/s15327752jpa4901_13
- Diener, E., Oishi, S., & Lucas, R. E. (2009). Subjective Well-Being: the science of happiness and life satisfaction. In *Handbook of positive psychology* (pp. 186–194). <https://doi.org/10.1093/oxfordhb/9780195187243.013.0017>

- Epstein, R. M., & Krasner, M. S. (2013). Physician resilience. *Academic Medicine*, *88*(3), 301–303. <https://doi.org/10.1097/acm.0b013e318280cff0>
- Fazekas, C., Zieser, M., Hanfstingl, B., Saretzki, J., Kunschitz, E., Zieser-Stelzhammer, L., Linder, D., & Matzer, F. (2024). Physician resilience and perceived quality of care among medical doctors with training in psychosomatic medicine during the COVID-19 pandemic: a quantitative and qualitative analysis. *BMC Health Services Research*, *24*(1). <https://doi.org/10.1186/s12913-024-10681-1>
- Ferreira, P., & Gomes, S. (2021). The Role of Resilience in Reducing Burnout: A Study with Healthcare Workers during the COVID-19 Pandemic. *Social Sciences*, *10*(9), 317. <https://doi.org/10.3390/socsci10090317>
- Filaj, E., & Sota, M. (2025). THE IMPACT OF BURNOUT ON THE QUALITY OF PATIENT CARE: A COMPARATIVE ANALYSIS BETWEEN PHYSICIANS AND NURSES IN EMOTIONALLY DEMANDING HOSPITAL SETTINGS. *KNOWLEDGE - International Journal*, *72*(1), 153–157.
- Fitzpatrick, B., Bloore, K., & Blake, N. (2019). Joy in work and reducing nurse burnout: from triple aim to quadruple aim. *AACN Advanced Critical Care*, *30*(2), 185–188. <https://doi.org/10.4037/aacnacc2019833>
- Freudenberger, H. J. (1974). Staff Burn-Out. *Journal of Social Issues*, *30*(1), 159–165. <https://doi.org/10.1111/j.1540-4560.1974.tb00706.x>
- García, G. M., & Calvo, J. C. A. (2011). Emotional exhaustion of nursing staff: influence of emotional annoyance and resilience. *International Nursing Review*, *59*(1), 101–107. <https://doi.org/10.1111/j.1466-7657.2011.00927.x>

- Gómez, M. a. L., Sabbath, E., Boden, L., Williams, J. A., Hopcia, K., Hashimoto, D., & Sorensen, G. (2019). Organizational and psychosocial working conditions and their relationship with mental health outcomes in Patient-Care workers. *Journal of Occupational and Environmental Medicine*, *61*(12), e480–e485. <https://doi.org/10.1097/jom.0000000000001736>
- Gonzalez, L., Duque, A., Molins, M., Alino, M., Molins, F., & Gresa, P. M. (2025). Resilience and social support as predictors of life satisfaction in healthy older adults: A moderation analysis. *medRxiv (Cold Spring Harbor Laboratory)*. <https://doi.org/10.1101/2025.05.24.25328287>
- Grossi, G., Perski, A., Osika, W., & Savic, I. (2015). Stress-related exhaustion disorder – clinical manifestation of burnout? A review of assessment methods, sleep impairments, cognitive disturbances, and neuro-biological and physiological changes in clinical burnout. *Scandinavian Journal of Psychology*, *56*(6), 626–636. <https://doi.org/10.1111/sjop.1225>
- Guo, Y., Luo, Y., Lam, L., Cross, W., Plummer, V., & Zhang, J. (2017). Burnout and its association with resilience in nurses: A cross-sectional study. *Journal of Clinical Nursing*, *27*(1–2), 441–449. <https://doi.org/10.1111/jocn.13952>
- Hamdan, M., Haddad, B. I., Alshrouf, M. A., Al-Ani, A., Alisi, M. S., Hammad, Y., Alelaumi, A., Hawamdeh, H. A., Halaweh, A. A., Alshabat, L., & Kawasmi, S. (2023). Burnout, grit and resilience among Jordanian orthopedic surgeons: a cross-sectional study. *BMC Medical Education*, *23*(1). <https://doi.org/10.1186/s12909-023-04572-y>

- Haraldseid-Driftland, C., Billett, S., Guise, V., Schibeveag, L., Alsvik, J. G., Fagerdal, B., Lyng, H. B., & Wiig, S. (2022). The role of collaborative learning in resilience in healthcare— a thematic qualitative meta-synthesis of resilience narratives. *BMC Health Services Research, 22*(1). <https://doi.org/10.1186/s12913-022-08451-y>
- Harry, E., Sinsky, C., Dyrbye, L. N., Makowski, M. S., Trockel, M., Tutty, M., Carlasare, L. E., West, C. P., & Shanafelt, T. D. (2020). Physician task load and the risk of burnout among US physicians in a national survey. *The Joint Commission Journal on Quality and Patient Safety, 47*(2), 76–85. <https://doi.org/10.1016/j.jcjq.2020.09.011>
- Hashmi, A. M., Ahmed, B., Nawal, A., & Asif, A. (2021). Phoenix from the Ashes: Physician Burnout in Pakistan. *Journal of Pakistan Psychiatric Society, 18*(03), 30–33. <https://doi.org/10.63050/jpps.18.03.130>
- Hobfoll, S. E. (2010). Conservation of Resources Theory: Its implication for stress, health, and Resilience. In *Oxford University Press eBooks*. <https://doi.org/10.1093/oxfordhb/9780195375343.013.0007>
- Hodkinson, A., Zhou, A., Johnson, J., Geraghty, K., Riley, R., Zhou, A., Panagopoulou, E., Chew-Graham, C. A., Peters, D., Esmail, A., & Panagioti, M. (2022). Associations of physician burnout with career engagement and quality of patient care: systematic review and meta-analysis. *BMJ, e070442*. <https://doi.org/10.1136/bmj-2022-070442>
- Hussain, S. S., Qazi, Q., & Bawar, S. (2022). Burnout in doctors working in tertiary care hospital in peshawar, pakistan. *Journal of Postgraduate Medical Institute*. <https://doi.org/10.54079/jpmi.36.4.3031>
- Iskander, M. (2018). Burnout, cognitive overload, and metacognition in medicine. *Medical Science Educator, 29*(1), 325–328. <https://doi.org/10.1007/s40670-018-00654-5>

- Jackson, J. L., Kuriyama, A., & Muramatsu, K. (2023). A model of burnout among healthcare professionals. *Journal of General Internal Medicine*, *39*(3), 373–376. <https://doi.org/10.1007/s11606-023-08514-8>
- Jackson, S. E., Schwab, R. L., & Schuler, R. S. (1986). Toward an understanding of the burnout phenomenon. *Journal of Applied Psychology*, *71*(4), 630–640. <https://doi.org/10.1037/0021-9010.71.4.630>
- Karagöz, Y., Durak, İ., & Mutlu, H. T. (2021). Psychological resilience and life satisfaction of healthcare professionals in Turkey: The Mediating role of fear of COVID. *Türk Psikolojik Danışma Ve Rehberlik Dergisi*, *11*(62), 403–417. <https://doi.org/10.17066/tpdrd.1001724>
- Karataş, Z., & Tagay, Ö. (2020). The relationships between resilience of the adults affected by the covid pandemic in Turkey and Covid-19 fear, meaning in life, life satisfaction, intolerance of uncertainty and hope. *Personality and Individual Differences*, *172*, 110592. <https://doi.org/10.1016/j.paid.2020.110592>
- Khalid, N. M. (2021). RESILIENCE, PERCEIVED SOCIAL SUPPORT, AND LIFE SATISFACTION AMONG MALAYSIAN COLLEGE STUDENTS. *Journal of Nusantara Studies (JONUS)*, *6*(2), 21–40. <https://doi.org/10.24200/jonus.vol6iss2pp21-40>
- Kim, E. S., Park, N., Sun, J. K., Smith, J., & Peterson, C. (2013). Life satisfaction and frequency of doctor visits. *Psychosomatic Medicine*, *76*(1), 86–93. <https://doi.org/10.1097/psy.0000000000000024>
- Koźtuniuk, A., Witczak, I., Młynarska, A., Czajor, K., & Uchmanowicz, I. (2021). Satisfaction with life, satisfaction with job, and the level of care rationing among Polish Nurses—A Cross-Sectional Study. *Frontiers in Psychology*, *12*. <https://doi.org/10.3389/fpsyg.2021.734789>

- Kristensen, T. S., Borritz, M., Villadsen, E., & Christensen, K. B. (2005). The Copenhagen Burnout Inventory: A new tool for the assessment of burnout. *Work & Stress*, *19*(3), 192–207. <https://doi.org/10.1080/02678370500297720>
- Lacy, B. E., & Chan, J. L. (2017). Physician burnout: the hidden health care crisis. *Clinical Gastroenterology and Hepatology*, *16*(3), 311–317. <https://doi.org/10.1016/j.cgh.2017.06.043>
- Łaskawiec-żuławińska, D., Grajek, M., Krupa-Kotara, K., Szlacheta, P., Karacan, H., Roszak, M., Łabuz-Rozsak, B., & Korzonek-Szlacheta, I. (2024). Burnout and Life Satisfaction among Healthcare Workers Related to the COVID-19 Pandemic (Silesia, Poland). *Behavioural Neurology*, *2024*, 1–7. <https://doi.org/10.1155/2024/9945392>
- Leitão, J., Pereira, D., & Gonçalves, Â. (2019). Quality of work life and organizational performance: workers' feelings of contributing, or not, to the organization's productivity. *International Journal of Environmental Research and Public Health*, *16*(20), 3803. <https://doi.org/10.3390/ijerph16203803>
- Lenzenweger, M. F. (2004). Authentic happiness: using the new positive psychology to realize your potential for lasting fulfillment. *American Journal of Psychiatry*, *161*(5), 936–937. <https://doi.org/10.1176/appi.ajp.161.5.936>
- Liaqat, N., Dar, S. H., & Waheed, K. (2019). Burn Out among the Pediatric Surgeons Of Pakistan. *Journal of the Pakistan Medical Association*, *0*, 1. <https://doi.org/10.5455/jpma.300916>

- Liu, N., Plouffe, R. A., Liu, J. J. W., Nouri, M. S., Saha, P., Gargala, D., Davis, B. D., Nazarov, A., & Richardson, J. D. (2024). Determinants of burnout in Canadian health care workers during the COVID-19 pandemic. *European Journal of Psychotraumatology*, *15*(1). <https://doi.org/10.1080/20008066.2024.2351782>
- Lo, D., Wu, F., Chan, M., Chu, R., & Li, D. (2018). A systematic review of burnout among doctors in China: a cultural perspective. *Asia Pacific Family Medicine*, *17*(1). <https://doi.org/10.1186/s12930-018-0040-3>
- Lombardo, P., Jones, W., Wang, L., Shen, X., & Goldner, E. M. (2018). The fundamental association between mental health and life satisfaction: results from successive waves of a Canadian national survey. *BMC Public Health*, *18*(1). <https://doi.org/10.1186/s12889-018-5235-x>
- Low, Z. X., Yeo, K. A., Sharma, V. K., Leung, G. K., McIntyre, R. S., Guerrero, A., Lu, B., Lam, C. C. S. F., Tran, B. X., Nguyen, L. H., Ho, C. S., Tam, W. W., & Ho, R. C. (2019). Prevalence of burnout in Medical and Surgical Residents: A Meta-Analysis. *International Journal of Environmental Research and Public Health*, *16*(9), 1479. <https://doi.org/10.3390/ijerph16091479>
- Lozano, J. M. G., Ramón, J. P. M., & Rodríguez, F. M. M. (2021). Doctors and Nurses: A Systematic review of the risk and protective factors in workplace violence and burnout. *International Journal of Environmental Research and Public Health*, *18*(6), 3280. <https://doi.org/10.3390/ijerph18063280>
- Lyubarova, R., Salman, L., & Rittenberg, E. (2023). Gender differences in physician burnout: driving factors and potential solutions. *The Permanente Journal*, *27*(2), 130–136. <https://doi.org/10.7812/tpp/23.023>

- Mahmood, J. I., Grotmol, K. S., Tesli, M., Moum, T., Andreassen, O., & Tyssen, R. (2019). Life satisfaction in Norwegian medical doctors: a 15-year longitudinal study of work-related predictors. *BMC Health Services Research*, *19*(1). <https://doi.org/10.1186/s12913-019-4599-7>
- Mansi, N., Verma, N. R. K., & Dabas, N. A. (2025). Comparative Analysis of Stress Prevalence among Married and Unmarried Female Doctors: A Comparative cross-sectional study. *Journal of Ayurveda and Integrated Medical Sciences*, *10*(4), 38–45. <https://doi.org/10.21760/jaims.10.4.7>
- Marshall, A. L., Dyrbye, L. N., Shanafelt, T. D., Sinsky, C. A., Satele, D., Trockel, M., Tutty, M., & West, C. P. (2020). Disparities in burnout and satisfaction with Work–Life integration in U.S. physicians by gender and practice setting. *Academic Medicine*, *95*(9), 1435–1443. <https://doi.org/10.1097/acm.00000000000003521>
- Marzo, R. R., Khaled, Y., ElSherif, M., Abdullah, M. S. a. M. B., Thew, H. Z., Chong, C., Soh, S. Y., Siau, C. S., Chauhan, S., & Lin, Y. (2022). Burnout, resilience and the quality of life among Malaysian healthcare workers during the COVID-19 pandemic. *Frontiers in Public Health*, *10*. <https://doi.org/10.3389/fpubh.2022.1021497>
- Maslach, C., Schaufeli, W. B., & Leiter, M. P. (2001). Job burnout. *Annual Review of Psychology*, *52*(1), 397–422. <https://doi.org/10.1146/annurev.psych.52.1.397>
- Mazhar, S. B., Gilani, S., Ain, Q. T., & Khan, S. (2019). High burn out among doctors working in a tertiary care hospital; a wakeup call. *PubMed*, *69*(3), 349–354. <https://pubmed.ncbi.nlm.nih.gov/30890826>

- McCain, R. S., McKinley, N., Dempster, M., Campbell, W. J., & Kirk, S. J. (2017). A study of the relationship between resilience, burnout and coping strategies in doctors. *Postgraduate Medical Journal*, *94*(1107), 43–47. <https://doi.org/10.1136/postgradmedj-2016-134683>
- Medeni, V., Medeni, İ., Altunay, G., Dikmen, A. U., & İlhan, M. N. (2025). Job satisfaction, life satisfaction, and associated factors among hospital nurses: a cross-sectional study in Türkiye. *Scientific Reports*, *15*(1). <https://doi.org/10.1038/s41598-025-85564-4>
- Merkin, R. (2023). Editorial: Employee life satisfaction: real world consequences. *Frontiers in Psychology*, *14*. <https://doi.org/10.3389/fpsyg.2023.1288877>
- Meynaar, I. A., Ottens, T., Zegers, M., Van Mol, M. M., & Van Der Horst, I. C. (2020). Burnout, resilience and work engagement among Dutch intensivists in the aftermath of the COVID-19 crisis: A nationwide survey. *Journal of Critical Care*, *62*, 1–5. <https://doi.org/10.1016/j.jcrc.2020.11.010>
- Misfin, E. W., Singh, M., & Phoolka, S. (2024). Impact of family and work conflicts on job satisfaction, life satisfaction and health of female teachers in Ethiopia: the moderating role of resilience. *Cogent Business & Management*, *11*(1). <https://doi.org/10.1080/23311975.2024.2430458>
- Misiotek, A., Gil-Monte, P. R., & Misiotek, H. (2015). Prevalence of burnout in Polish anesthesiologists and anesthetist nursing professionals: A comparative non-randomized cross-sectional study. *Journal of Health Psychology*, *22*(4), 465–474. <https://doi.org/10.1177/1359105315604377>

- Moss, M., Good, V. S., Gozal, D., Kleinpell, R., & Sessler, C. N. (2016). An official Critical Care Societies Collaborative statement: Burnout Syndrome in Critical care Health care Professionals: A Call for Action. *American Journal of Critical Care*, 25(4), 368–376. <https://doi.org/10.4037/ajcc2016133>
- MSEd, K. C. (2023, May 4). *Subjective Well-Being—A way to measure your own happiness*. Verywell Mind. <https://www.verywellmind.com/what-is-subjective-well-being-5221255>
- Murtaza, G. (2025). Burnout among psychiatry doctors in Pakistan: a multicenter study. *European Psychiatry*, 68(S1), S798. <https://doi.org/10.1192/j.eurpsy.2025.1622>
- Nijimbere, A., & Hussain, I. (2017). Academic stress and resilience among married and unmarried students at Hope Africa University, Burundi. *International Journal of Economic Research*, 14(20), 811-816.
- Nituica, C., Bota, O. A., & Blebea, J. (2020). Specialty differences in resident resilience and burnout - A national survey. *The American Journal of Surgery*, 222(2), 319–328. <https://doi.org/10.1016/j.amjsurg.2020.12.039>
- Otukoya, E. Z., Amiri, A., & Alimohammadi, E. (2025). Surgeon well-being: a systematic review of stressors, mental health, and resilience. *BMC Surgery*, 25(1). <https://doi.org/10.1186/s12893-025-03180-5>
- Pastores, S. M., Kvetan, V., Coopersmith, C. M., Farmer, J. C., Sessler, C., Christman, J. W., D'Agostino, R., Diaz-Gomez, J., Gregg, S. R., Khan, R. A., Kapu, A. N., Masur, H., Mehta, G., Moore, J., Oropello, J. M., & Price, K. (2019). Workforce, Workload, and burnout among intensivists and advanced practice Providers: A Narrative review. *Critical Care Medicine*, 47(4), 550–557. <https://doi.org/10.1097/ccm.0000000000003637>

- Permarupan, P. Y., Mamun, A. A., Samy, N. K., Saufi, R. A., & Hayat, N. (2020). Predicting Nurses Burnout through Quality of Work Life and Psychological Empowerment: A Study Towards Sustainable Healthcare Services in Malaysia. *Sustainability*, *12*(1), 388. <https://doi.org/10.3390/su12010388>
- Petrakova, A. (n.d.). Exploring the evolution of psychological well-being concept. *www.alliedacademies.org*. <https://doi.org/10.35841/aajmha-7.1.132>
- Phillips, G., Cameron, A., & Szakmany, T. (2019). Defining the high-risk patient. *Journal of Emergency and Critical Care Medicine*, *3*, 40. <https://doi.org/10.21037/jeccm.2019.08.10>
- Robertson, H. D., Elliott, A. M., Burton, C., Iversen, L., Murchie, P., Porteous, T., & Matheson, C. (2016). Resilience of primary healthcare professionals: a systematic review. *British Journal of General Practice*, *66*(647), e423–e433. <https://doi.org/10.3399/bjgp16x685261>
- Romani, M., & Ashkar, K. (2014). Burnout among physicians. *Libyan Journal of Medicine*, *9*(1). <https://doi.org/10.3402/ljm.v9.23556>
- Rotenstein, L., Harry, E., Wickner, P., Gupte, A., Neville, B. A., Lipsitz, S., Cullen, E., Rozenblum, R., Sequist, T. D., & Dudley, J. (2021). Contributors to gender Differences in burnout and Professional Fulfillment: A survey of physician faculty. *The Joint Commission Journal on Quality and Patient Safety*, *47*(11), 723–730. <https://doi.org/10.1016/j.jcjq.2021.08.002>
- Rushton, C. H., Batcheller, J., Schroeder, K., & Donohue, P. (2015). Burnout and resilience among nurses practicing in High-Intensity Settings. *American Journal of Critical Care*, *24*(5), 412–420. <https://doi.org/10.4037/ajcc2015291>

- Sanfilippo, F., Noto, A., Palumbo, G. J., Ippolito, M., Gagliardone, M., Scarlata, M., Bignami, E., Sangalli, F., Cattaneo, S., Blangetti, I., Scolletta, S., Locatelli, A., Tritapepe, L., Lorini, F. L., & Arcadipane, A. (2018). Burnout in cardiac anesthesiologists: results from a national survey in Italy. *Journal of Cardiothoracic and Vascular Anesthesia*, *32*(6), 2459–2466. <https://doi.org/10.1053/j.jvca.2018.05.016>
- Sarwar, S., Mahmood, A., Raja, M. F., & Mahmud, Y. (2023). Burnout and Health Related Quality of Life among emergency physicians working at emergency at Tertiary Care Hospital in Lahore Pakistan. *Pakistan Journal of Medical Sciences*, *39*(5). <https://doi.org/10.12669/pjms.39.5.7560>
- Sclanavo, M. (2023). Examining Resilience and its Role in Positive Psychological Outcomes. *Interdisciplinary Journal Papier Human Review*, *4*(1), 1–9. <https://doi.org/10.47667/ijphr.v4i1.233>
- Shanafelt, T. D., Balch, C. M., Bechamps, G. J., Russell, T., Dyrbye, L., Satele, D., Collicott, P., Novotny, P. J., Sloan, J., & Freischlag, J. A. (2009). Burnout and career satisfaction among American surgeons. *Annals of Surgery*, *250*(3), 463–471. <https://doi.org/10.1097/sla.0b013e3181ac4dfd>
- Sheikhrabori, A., Peyrovi, H., & Khankeh, H. (2022). The main features of resilience in healthcare providers: A scoping review. *Medical Journal of the Islamic Republic of Iran*. <https://doi.org/10.47176/mjiri.36.3>
- Shi, M., Wang, X., Bian, Y., & Wang, L. (2015). The mediating role of resilience in the relationship between stress and life satisfaction among Chinese medical students: a cross-sectional study. *BMC Medical Education*, *15*(1). <https://doi.org/10.1186/s12909-015-0297-2>

- Shim, T., & Go, E. (2025). The impact of school burnout on life satisfaction among university students: The mediating effects of loneliness and fear of alienation. *Behavioral Sciences, 15*(8), 1083. <https://doi.org/10.3390/bs15081083>
- Sinbukhova, E. V., Petrikov, S. S., Shabanov, A. K., Plotnikov, G. P., Lubnin, A. Y., & Sinbukhova, N. I. (2019). Open Questions of Life Satisfaction and Burnout in Anesthesiology and Resuscitation. *Russian Sklifosovsky Journal Emergency Medical Care, 8*(3), 266–273. <https://doi.org/10.23934/2223-9022-2019-8-3-266-273>
- Smith, B. W., Dalen, J., Wiggins, K., Tooley, E., Christopher, P., & Bernard, J. (2008). The brief resilience scale: Assessing the ability to bounce back. *International Journal of Behavioral Medicine, 15*(3), 194–200. <https://doi.org/10.1080/10705500802222972>
- Soler, J. K., Yaman, H., Esteva, M., Dobbs, F., Asenova, R. S., Katic, M., Ozvacic, Z., Desgranges, J. P., Moreau, A., Lionis, C., Kotanyi, P., Carelli, F., Nowak, P. R., De Aguiar Sa Azeredo, Z., Marklund, E., Churchill, D., & Ungan, M. (2008). Burnout in European family doctors: the EGPRN study. *Family Practice, 25*(4), 245–265. <https://doi.org/10.1093/fampra/cmn038>
- Sultana, M., Sharmin, N., Karim, M. Z., & Sarkar, N. (2024). Resilience as predictor of life satisfaction and mental Well-Being among selected Southern University students of Bangladesh. *Bangladesh Psychological Association, 24*(1), 49–60. <https://doi.org/10.3329/bjop.v24i1.78523>
- Uchmanowicz, I., Manulik, S., Lomper, K., Rozensztrauch, A., Zborowska, A., Kolasińska, J., & Rosińczuk, J. (2019). Life satisfaction, job satisfaction, life orientation and occupational burnout among nurses and midwives in medical institutions in Poland: a cross-sectional study. *BMJ Open, 9*(1), e024296. <https://doi.org/10.1136/bmjopen-2018-024296>

- Vella, S., & Pai, N. (2019). A theoretical review of psychological resilience: Defining resilience and resilience research over the decades. *Archives of Medicine and Health Sciences*, 7(2), 233. https://doi.org/10.4103/amhs.amhs_119_19
- Vittersø, J. (2025). Life satisfaction. In *Humanistic Wellbeing (pp.145-182)* (pp. 145–182). https://doi.org/10.1007/978-3-031-69292-5_4
- Wagnild, G. M., & Collins, J. A. (2009). Assessing resilience. *Journal of Psychosocial Nursing and Mental Health Services*, 47(12), 28–33. <https://doi.org/10.3928/02793695-20091103-01>
- Wang, Q., Sun, W., & Wu, H. (2022). Associations between academic burnout, resilience and life satisfaction among medical students: a three-wave longitudinal study. *BMC Medical Education*, 22(1). <https://doi.org/10.1186/s12909-022-03326-6>
- Wang, W., Li, M., Zhang, J., Zhao, R., Yang, H., & Mitchell, R. (2024). Organizational resilience and primary care nurses' work conditions and wellbeing: a multilevel empirical study in China. *Health Policy and Planning*. <https://doi.org/10.1093/heapol/czae091>
- Werner, E. (1997). Vulnerable but invincible: high-risk children from birth to adulthood. *Acta Paediatrica*, 86(S422), 103–105. <https://doi.org/10.1111/j.1651-2227.1997.tb18356.x>
- West, C. P., Dyrbye, L. N., Sinsky, C., Trockel, M., Tutty, M., Nedelec, L., Carlasare, L. E., & Shanafelt, T. D. (2020). Resilience and burnout among physicians and the general US working population. *JAMA Network Open*, 3(7), e209385. <https://doi.org/10.1001/jamanetworkopen.2020.9385>

- Wijayatunga, S. L. (2021). Life Satisfaction among Healthcare Workers: A Cross-Sectional Study in Private Hospitals in Ragama, Sri Lanka » *The International Journal of Indian Psychology*. *International Journal of Indian Psychology*. <https://doi.org/10.25215/0903.185>
- World Health Organization: WHO. (2019, May 28). *Burn-out an "occupational phenomenon": International Classification of Diseases*. <https://www.who.int/news/item/28-05-2019-burn-out-an-occupational-phenomenon-international-classification-of-diseases>
- Wu, N., Ding, F., Zhang, R., Cai, Y., & Zhang, H. (2022). The Relationship between Perceived Social Support and Life Satisfaction: The Chain Mediating Effect of Resilience and Depression among Chinese Medical Staff. *International Journal of Environmental Research and Public Health*, *19*(24), 16646. <https://doi.org/10.3390/ijerph192416646>
- Wu, S., Zhu, W., Wang, Z., Wang, M., & Lan, Y. (2007). Relationship between burnout and occupational stress among nurses in China. *Journal of Advanced Nursing*, *59*(3), 233–239. <https://doi.org/10.1111/j.1365-2648.2007.04301.x>
- Zahednezhad, H., Zareiyan, A., & Jame, S. Z. B. (2021). Relationship between quality of work-life, resilience and burnout among nursing professionals during COVID-19 pandemic in Iran: A cross-sectional study. *Belitung Nursing Journal*, *7*(6), 508–515. <https://doi.org/10.33546/bnj.1702>
- Załuski, M., & Makara-Studzińska, M. (2024). Having children and being married are predictors of burnout and obesity among working men: Effects of Latent Profile analysis. *INQUIRY the Journal of Health Care Organization Provision and Financing*, *61*, 469580241229642. <https://doi.org/10.1177/00469580241229642>

Zhang, Q., Mu, M., He, Y., Cai, Z., & Li, Z. (2020). Burnout in emergency medicine physicians. *Medicine*, 99(32), e21462. <https://doi.org/10.1097/md.00000000000021462>