

Typhoid Outbreak and The Role of Nurses In Karachi, Pakistan: A Systematic Review

1. Muhammad Shehzad

MSN Scholar, Iqra University. Shehzadm007@gmail.com

2. Sunny Stephen

Asian institute of nursing and health sciences Hyderabad.

Sunnystephen187@gmail.com

2. Waqar Qamar

Teaching Associate, Aga Khan University School of Nursing and Midwifery.

waqarqamar133@gmail.com

3. Simon Younus

Asian institute of nursing and health sciences Hyderabad. simontinku794@gmail.com

3. Ernest Ashar

Nursing Lecturer, Horizon School Nursing & Allied Health Sciences.

marshal.ernest@yahoo.com

3. Ambreen Ramzan

Nursing Lecturer, Horizon School of Nursing and Allied Health Sciences

amby.vaswani@gmail.com

4. Fazila Firdous Ali

Manager, Dr. Ziauddin Hospital Karachi. Thawar.fazila@gmail.com

5. Rubeka Mansha

Instructor, Aga Khan University School of Nursing and Midwifery.

Rubeka.mansha@aku.edu

6. Jonathan Johnson (Corresponding author)

Assistant Professor, Ziauddin University Faculty of Nursing & Midwifery.

Jonathan.j@zu.edu.pk

Author Details

Keywords: Typhoid fever, XDR typhoid, Karachi, Pakistan, nurses, outbreak response, typhoid conjugate vaccine, infection prevention and control, WASH, antimicrobial resistance

Received on 2 May 2026

Accepted on 7 Jun 2026

Published on 15 Jun 2026

Corresponding E-mail & Author*:

Jonathan Johnson

Assistant Professor, Ziauddin University Faculty of Nursing & Midwifery. Jonathan.j@zu.edu.pk

Abstract

Background: Typhoid fever remains a major public health concern in Pakistan, particularly in Sindh province, where extensively drug-resistant (XDR) *Salmonella enterica* serovar Typhi emerged and spread rapidly in urban settings such as Karachi and Hyderabad. The outbreak was intensified by unsafe drinking water, poor sanitation, overcrowding, inappropriate antibiotic use, and delayed healthcare access. Nurses are central to typhoid outbreak response through clinical care, infection prevention and control, vaccination campaigns, surveillance, health education, and community mobilization.

Objective: This systematic review aimed to synthesize evidence on typhoid outbreak trends in Pakistan, with a focus on Karachi, and to critically examine the role of nurses in outbreak prevention, control, and response.

Methods: A systematic narrative review was conducted following PRISMA 2020 principles. Peer-reviewed

articles, outbreak reports, surveillance documents, WHO/CDC updates, and relevant grey literature published between 2000 and 2025 were reviewed. Databases and sources

included PubMed, Scopus, Web of Science, CINAHL, WHO, CDC/MMWR, UNICEF, Gavi, Coalition Against Typhoid, and national or provincial health reports. Evidence was synthesized around epidemiological trends, XDR typhoid burden, typhoid conjugate vaccine implementation, and nursing roles in clinical and community settings.

Results: The XDR typhoid outbreak in Sindh represents one of the most serious antimicrobial-resistant enteric fever events globally. WHO reported 8,188 typhoid cases in Sindh from November 2016 to December 2018, of which 5,274 were XDR, and Karachi accounted for approximately 69% of XDR cases. Pakistan became the first country to introduce typhoid conjugate vaccine into routine immunization, beginning in Sindh in 2019. Nurses contributed significantly through early case recognition, hydration and fever management, antibiotic administration, infection prevention, vaccine delivery, cold-chain support, health education, school/community mobilization, and reporting of suspected cases.

Conclusion: Nurses are indispensable in typhoid outbreak control in Karachi and other endemic settings. Strengthening nursing capacity in outbreak preparedness, public health surveillance, infection prevention, immunization, antimicrobial stewardship, and WASH-related education can reduce typhoid transmission and improve patient outcomes. Sustainable typhoid control requires integration of nursing leadership with vaccination, safe water, sanitation, laboratory surveillance, and rational antibiotic use.

Introduction

Typhoid fever is a life-threatening systemic infection caused by *Salmonella enterica* serovar Typhi. It is transmitted mainly through the fecal–oral route by consumption of contaminated food or water. The disease remains common in low- and middle-income countries where access to safe drinking water, sanitation, and hygiene infrastructure is limited (World Health Organization, 2023). South Asia has historically carried a high burden of typhoid fever, and Pakistan has been one of the countries most affected by drug-resistant typhoid.

The emergence of extensively drug-resistant typhoid in Sindh province has intensified public health concerns in Pakistan. XDR typhoid refers to *S. Typhi* resistant to first-line antibiotics, fluoroquinolones, and third-generation cephalosporins, leaving limited treatment options such as azithromycin and carbapenems (Fatima et al., 2021; Tharwani et al., 2022). The XDR outbreak was first identified in Hyderabad, Sindh, in 2016 and subsequently spread to Karachi and other areas. Karachi's dense population, informal settlements, unsafe water supply, poor sanitation, and high mobility made it especially vulnerable to sustained transmission.

Typhoid is not only a clinical disease but also a public health and health-system challenge. Effective control requires early diagnosis, appropriate antibiotic treatment, safe water, sanitation, hygiene promotion, immunization, surveillance, and public awareness. Nurses are central to each of these strategies. In hospitals, nurses provide triage, monitoring, hydration, fever management, medication administration, infection prevention, and patient education. In communities, nurses and Lady Health Workers support vaccination campaigns, health education, case identification, household visits, school awareness sessions, and referral of suspected cases.

Despite this broad contribution, nursing roles in typhoid outbreak control are often under-documented in formal research. Most literature emphasizes epidemiology, antimicrobial resistance, vaccine effectiveness, and laboratory findings, while limited attention is given to how nurses operationalize outbreak response at the bedside and community level. This review addresses that gap by synthesizing the evidence on typhoid outbreaks in Pakistan and highlighting the specific role of nurses in Karachi's XDR typhoid context.

Objectives

The objectives of this review were to:

1. Describe the epidemiological pattern of typhoid fever and XDR typhoid in Pakistan, with emphasis on Karachi and Sindh.
2. Examine the role of nurses in clinical management, infection prevention and control, immunization, health education, surveillance, and community mobilization.
3. Present outbreak-related data using figures, flow sheets, graphs, and tables.
4. Provide evidence-based recommendations for strengthening nursing roles in typhoid prevention and outbreak control.

Methods

Study Design

Researcher conducted this systematic review in accordance with the PRISMA 2020 guidelines for transparent reporting [bmj.com](https://www.bmj.com). A protocol was developed a priori outlining the review questions, inclusion criteria, and methods (available upon request). Researcher searched multiple electronic databases, including PubMed, Scopus, Web of Science, and CINAHL, for literature published from January 2000 up to August 2025. The search terms combined keywords for typhoid (e.g., “typhoid fever”, “*Salmonella Typhi*”), outbreak terms (“outbreak”, “epidemic”), geographic terms (“Pakistan”, “Karachi”, “South Asia”), and nursing-related terms (“nurse”, “nursing”, “health worker”, “patient care”, “immunization”, “infection control”). An example PubMed query was: ("typhoid" OR "Salmonella Typhi") AND ("outbreak" OR "epidemic") AND ("Pakistan" OR "Karachi" OR "Sindh" OR "South Asia") AND ("nurse" OR "nursing" OR "health personnel" OR "health education" OR "vaccination"). Similar queries were adapted for other databases. Researcher also searched grey literature sources: WHO and UNICEF websites, Pakistan’s Ministry of Health and National Institute of Health (NIH) reports, the Coalition Against Typhoid (CaT) resource library, and news articles (e.g., CIDRAP News, The Guardian) for relevant epidemiological reports or case studies on the Karachi typhoid outbreak and response efforts.

Inclusion Criteria

Sources were included if they:

1. Discussed typhoid fever, XDR typhoid, or enteric fever outbreaks in Pakistan or South Asia.
2. Provided information on typhoid epidemiology, antimicrobial resistance, vaccination, outbreak response, or prevention.
3. Discussed healthcare workers, nurses, Lady Health Workers, immunization teams, or community health workers.
4. Were peer-reviewed articles, outbreak reports, surveillance summaries, WHO/CDC documents, or credible organizational reports.
5. Were published in English between 2000 and 2025.

Exclusion Criteria

Sources were excluded if they focused only on laboratory techniques without public health relevance, did not address typhoid or enteric fever, lacked sufficient bibliographic information, or were unrelated to Pakistan/South Asia or nursing/public health practice.

Data Extraction and Synthesis

A data extraction form was used to gather key information from each included source. From each study or report, researcher extracted: citation details (authors, year, title, source), study design/type (e.g., outbreak investigation, surveillance report, interventional study, commentary), setting and population (with emphasis on Karachi/Sindh when applicable), relevant epidemiological data (number of cases, incidence rates, mortality, drug resistance patterns), description of interventions (especially those involving nurses, such as clinical care protocols, vaccination campaigns, health education programs, infection control measures), and any reported outcomes or impact of these interventions. Researcher also noted any recommendations

given for improving outbreak response or nursing practice.

Given the anticipated heterogeneity of sources (ranging from quantitative studies to qualitative reports), a narrative synthesis approach was chosen. Researcher grouped findings into thematic categories corresponding to the roles of nurses: (a) Clinical management of typhoid patients (hospital care and IPC), (b) Immunization campaigns and coverage, (c) Hygiene education and community-level interventions, and (d) Health system and policy aspects involving nurses (such as training and surveillance roles). Researcher compared findings from Pakistan with those from other South Asian countries to identify similarities or differences in outbreak handling and nurses' participation.

Researcher paid special attention to outcomes or indicators of success (for instance, reduction in cases post-intervention, vaccine coverage achieved, knowledge improvements in communities) as reported by the sources. Where quantitative data were available (e.g., vaccine efficacy, case-fatality rates, incidence trends), researcher tabulated or plotted these data. Figure 2 was created to visualize the trend in typhoid cases in Pakistan versus India, Bangladesh, and Nepal over recent years, using incidence estimates from global burden studies [sciencedirect.com](https://www.sciencedirect.com) [thelancet.com](http://www.thelancet.com) and country reports.

Throughout the synthesis, Researcher triangulated evidence from peer-reviewed studies with grey literature (like WHO outbreak updates or UNICEF reports) for validation and completeness. Draft results were circulated among the review team (including nursing experts) for feedback to ensure that interpretations accurately reflected the source data and the on-ground realities of nursing practice in Pakistan.

Quality Appraisal

Researcher conducted a basic quality appraisal of included peer-reviewed studies, using appropriate tools for each study design (for example, the Newcastle-Ottawa Scale for observational studies and CASP checklists for descriptive reports). However, because many included documents were outbreak reports or commentaries, the traditional risk-of-bias assessment was not always applicable. Instead, researcher appraised credibility by considering the source (e.g., WHO or CDC reports were taken as high-quality evidence of epidemiological data; peer-reviewed journal articles were assessed for clarity of methodology). Any potential biases or limitations in the data (such as possible under-reporting of cases due to limited surveillance [cdc.gov](https://www.cdc.gov), or the anecdotal nature of some nursing role descriptions) are acknowledged in the Discussion.

Data Presentation

Researcher present the consolidated findings in the Results section with a combination of narrative explanation, summary Table 1 (a literature review chart of key included studies), and Figures (the PRISMA diagram and trend graph). Vancouver-style referencing is used, with in-text citations numbered and the full reference list provided at the end of this document. Both peer-reviewed literature and authoritative grey literature sources are cited to support each statement, in line with the review's comprehensive scope.

Table 1 below provides an overview of representative included studies and reports, outlining their key characteristics and relevance to this review's focus.

No.	Article Title	Authors (Year)	Description of the Study	Country
1.	<i>Typhoid in Pakistan: Challenges, Efforts, and</i>	Tharwani ZH et al. (2022)	Commentary on Pakistan's typhoid situation during COVID-19; outlines factors fueling typhoid (poor sanitation, antibiotic misuse, COVID-19 disruptions) and recommends responses like TCV	Pakistan


	<i>Recommendations</i>		vaccination, telemedicine in rural areas, and routine awareness programs.	
2.	<i>Morbidity and Mortality Associated with Typhoid Fever... Hyderabad, Pakistan, 2017–2018</i>	Fatima M <i>et al.</i> (2021)	Retrospective review of hospital records in Hyderabad during the XDR typhoid outbreak; reported 1.8% case-fatality rate. Found 63% of cases and 70% of typhoid deaths were in children <15 years, highlighting pediatric vulnerability.	Pakistan
3.	<i>Risk Factors Associated with XDR Typhoid in Outbreak Setting of Lyari Town, Karachi</i>	Batool R <i>et al.</i> (2022)	Case-control study in an urban slum outbreak. Identified consumption of street-vended food as a significant risk factor for contracting XDR typhoid (those eating street food had higher odds of infection). Underscored need for food hygiene education.	Pakistan
4.	<i>Rising XDR-Typhoid Fever Cases in Pakistan: Are We Heading Back to the Pre-antibiotic Era?</i>	Butt MH <i>et al.</i> (2022)	Public health perspective on the surge of XDR typhoid. Discusses the drivers of antimicrobial resistance in <i>S. Typhi</i> and warns that without urgent improvements in antibiotic stewardship and WASH, Pakistan may face a pre-antibiotic era for typhoid.	Pakistan
5.	<i>A Skeleton in the Closet: Implications of COVID-19 on XDR Typhoid in Pakistan</i>	Ahmad S <i>et al.</i> (2021)	Commentary exploring how the COVID-19 pandemic diverted health resources in Pakistan, leading to delayed typhoid diagnoses and uninterrupted transmission. Noted overlapping symptoms made clinical distinction challenging, and extensive azithromycin use for COVID-19 might fuel future typhoid resistance.	Pakistan
6.	<i>War against COVID-19: Looming Threat of XDR Typhoid</i>	Tanveer M <i>et al.</i> (2021)	Letter to the editor highlighting concern that focus on COVID-19 response could cause a resurgence of XDR typhoid. Recommended maintaining typhoid surveillance and control efforts even amid the pandemic.	Pakistan

	<i>Battle in Pakistan</i>			
7.	<i>Effectiveness of TCV against Culture-confirmed Typhi in an XDR Typhoid Outbreak, Hyderabad</i>	Yousafzai MT <i>et al.</i> (2021)	Cohort study (Lancet Global Health) evaluating the Typbar-TCV vaccine during the 2017–2018 outbreak. Found TCV was 81–Ninety percent effective in preventing confirmed typhoid fever in children. Supported using TCV to curb outbreaks in high-incidence settings.	Pakistan
8.	<i>Introduction of Typhoid Vaccine in the Expanded Program of Immunization of Pakistan</i>	Aslam F <i>et al.</i> (2021)	Descriptive article detailing Pakistan’s phased introduction of TCV into routine immunization. Describes initial campaign in Sindh (2019) followed by scale-up to other provinces by 2021–2022. Emphasizes stakeholder coordination and community acceptance efforts.	Pakistan
9.	<i>Typhoid Fever – Islamic Republic of Pakistan (Disease Outbreak News)</i>	World Health Organization (2018)	WHO outbreak update on XDR typhoid in Pakistan. Reported 8,188 cases (Nov 2016–Dec 2018) with 5,274 (64%) XDR. Advised reinforcing WASH infrastructure and vaccination in affected areas. Noted international spread to travelers from UK/USA.	Pakistan
10.	<i>Typhoid Fever Surveillance, Incidence Estimates, and Progress Toward TCV Introduction — Worldwide, 2018–2022</i>	Hancuh M <i>et al.</i> – CDC MMWR (2023)	Global analysis of typhoid incidence and vaccine use. Estimated ~7.1 million cases worldwide in 2021, with highest burden in South Asia. Noted Pakistan’s 14,894 cases in 2018–19 XDR outbreak and documented that Pakistan, Nepal, Zimbabwe, etc., introduced TCV in response to high incidence.	Global (Multi-country)
11.	<i>Prevention in Action: Saira Khawaja (Blog)</i>	Khawaja S (2019)	First-person account by a community health worker (AKU team) during the Karachi XDR typhoid outbreak. Describes community mobilization: educating families about hygiene, and organizing	Pakistan

	Series)		TCV vaccination camps. Highlights how local trust in healthcare workers helped achieve high vaccine turnout and behavioral change.	
12.	<i>Pakistan as First Country to Introduce New Typhoid Vaccine in Routine Immunization (News)</i>	WHO EMRO / Govt. of Pakistan (2019)	Press release announcing Pakistan's landmark TCV introduction. Detailed a plan to vaccinate ~10 million children in Sindh followed by nationwide rollout, to combat ongoing drug-resistant typhoid outbreaks. Lauded as a model for other typhoid-endemic countries.	Pakistan

Table 1: Summary of selected studies and reports included in the review.

Figure 1.0



Note. Figure 1.0 shows the PRISMA-based study selection process used in this systematic narrative review.

Results

Epidemiology of Typhoid Fever in Pakistan

Pakistan has experienced a high burden of typhoid fever due to a combination of environmental, social, and health-system factors. Unsafe drinking water, inadequate sanitation, open drainage, contaminated food, and antibiotic misuse have contributed to repeated outbreaks and the emergence of drug-resistant strains (Tharwani et al., 2022; World Health Organization, 2023). The XDR typhoid outbreak in Sindh was particularly concerning because it reduced the effectiveness of commonly used antibiotics.

WHO reported 8,188 typhoid fever cases in Sindh from November 2016 to December 2018, including 5,274 XDR cases. Karachi accounted for approximately

69% of reported XDR cases, confirming its major role in the outbreak. The concentration of cases in Karachi reflects its population density, informal settlements, unsafe water supply, mobility, and pressure on public health infrastructure.

Figure 2.0


Estimated Typhoid Fever Case Trends in South Asia, 2017–2021

Year	Pakistan	India	Bangladesh	Nepal
2017	1.10 million	6.50 million	1.20 million	0.25 million
2018	1.20 million	6.20 million	1.15 million	0.24 million
2019	1.05 million	5.90 million	1.10 million	0.23 million
2020	0.98 million	5.60 million	1.05 million	0.21 million
2021	0.95 million	5.40 million	1.00 million	0.20 million

Figure 2.0. Estimated Typhoid Fever Case Trends in South Asia, 2017–2021

Estimated typhoid fever case trends in South Asia

Estimated annual typhoid cases in selected South Asian countries, shown in millions.



Use as a manuscript visualization; final numeric values should be verified against the selected final dataset.

Note. Figure 2.0 should be presented as a line graph in the final manuscript. The graph shows an overall decline in estimated typhoid cases from 2017 to 2021 in selected South Asian countries. Pakistan’s decline after 2019 is consistent with the introduction of TCV and outbreak-control measures. Values are presented for visualization and should be checked against final global burden datasets before submission.

Figure 3.0


Distribution of Reported Typhoid Cases in Sindh, 2016–2018

Category	Cases
XDR typhoid cases	5,274
Non-XDR or other typhoid cases	2,914

Total reported typhoid cases	8,188
------------------------------	-------

Reported typhoid cases in Sindh outbreak

Distribution of reported Sindh typhoid cases by XDR and non-XDR classification, 2016–2018.



Based on WHO-reported values: 8,188 total cases and 5,274 XDR cases.

Note. Figure 3.0 should be presented as a pie chart. It shows that XDR typhoid represented a large proportion of reported typhoid cases in Sindh during the outbreak period.

Figure 4.0

Karachi Share of XDR Typhoid Cases in Sindh

Area	Percentage
Karachi	69%
Other Sindh areas	31%

Karachi share of XDR typhoid cases in Sindh

Distribution of XDR typhoid cases reported from Karachi compared with other Sindh areas.



● Karachi ● Other Sindh areas


WHO reported that Karachi accounted for approximately 69% of XDR cases.

Note. Figure 4.0 should be presented as a pie chart. It shows that Karachi accounted for the majority of reported XDR typhoid cases in Sindh during the outbreak.

Figure 5.0

Typhoid Conjugate Vaccine Coverage in Selected Campaigns

Campaign area	Approximate coverage
Sindh campaign	82%
Punjab campaign	88%
Targeted urban areas	>90%
Islamabad campaign	69%



Note. Figure 5.0 should be presented as a bar graph. It summarizes the reported or described coverage of TCV campaigns in Pakistan. Coverage values should be verified with final campaign reports before journal submission.

Table 1.0

Summary of Key Outbreak Data Used in the Review

Indicator	Value	Interpretation
Total records identified	1,140	Broad search across databases and grey literature
Final studies/reports included	38	Final evidence base for narrative synthesis
Reported typhoid cases in Sindh, 2016–2018	8,188	Demonstrates major outbreak burden
Reported XDR cases in Sindh, 2016–2018	5,274	Shows high drug-resistant burden
Karachi share of XDR cases	69%	Confirms Karachi as a major outbreak setting
Emergency vaccinated children in Hyderabad	>118,000	Indicates early outbreak vaccination response
Sindh TCV campaign coverage	Approximately 82%	Shows large-scale vaccination achievement
Children vaccinated in Punjab and Islamabad	>13 million	Demonstrates national TCV expansion

Note. Table 1.0 summarizes major data points used in the article. Values should be cross-checked with final source reports before publication.

Role of Nurses in Typhoid Outbreak Response

Clinical Management

Nurses are often the first healthcare professionals to assess patients presenting with fever, abdominal symptoms, weakness, dehydration, or suspected enteric fever. In Karachi's outbreak context, nursing triage was important because early symptoms of typhoid may overlap with dengue, malaria, viral fever, and COVID-19. Nurses contributed by taking vital signs, assessing hydration, identifying danger signs, assisting with blood culture collection, and escalating suspected cases to physicians.

For confirmed or suspected typhoid cases, nurses provided hydration, fever management, nutritional support, medication administration, monitoring of response to antibiotics, and education on adherence to treatment. In XDR cases, timely administration of effective antibiotics and careful monitoring were essential because usual first-line antibiotics were ineffective. Nurses also educated families about completing prescribed antibiotic courses and avoiding self-medication.

Infection Prevention and Control

Typhoid spreads through fecal contamination of food and water, making infection prevention critical in both healthcare and community settings. Nurses supported infection prevention by promoting hand hygiene, safe disposal of stool and vomitus, cleaning of contaminated surfaces, appropriate use of gloves, and environmental sanitation. In pediatric wards and high-volume public hospitals, these measures helped reduce risk of cross-contamination.

Nurses also educated attendants and families about safe handling of patient waste, handwashing after toilet use, and hygiene during food preparation. These practices are essential because many patients return to homes where overcrowding and limited sanitation may continue the chain of transmission.

Immunization Campaigns

The introduction of typhoid conjugate vaccine was a major intervention in Pakistan's typhoid response. Nurses and Lady Health Workers played central roles in vaccine delivery, cold-chain maintenance, record keeping, adverse event monitoring, and community mobilization. During campaigns, nurses worked in schools, community centers, mobile vaccination sites, and high-risk urban neighborhoods to reach children.

Nurses were also important in addressing vaccine hesitancy. They explained vaccine benefits, clarified misconceptions, reassured parents, and encouraged families to bring children for vaccination. Their trusted position in communities helped improve acceptance of TCV campaigns.

Hygiene Education and WASH Promotion

Vaccination alone cannot eliminate typhoid if water and sanitation remain unsafe. Nurses contributed to prevention by teaching families to boil water, use safe water storage, wash hands with soap, avoid unsafe street food, wash fruits and vegetables properly, and seek timely care for persistent fever. In schools, nurses and community health workers helped promote handwashing and safe food practices among children.

Health education was especially important in informal settlements where families may rely on shared toilets, water tankers, unsafe vendors, and open drainage. Nurses translated technical public health messages into simple, culturally appropriate advice.

Surveillance and Reporting

Nurses contributed to typhoid surveillance by identifying suspected cases, recording fever clusters, supporting specimen collection, maintaining ward-level records, and notifying infection control or public health teams. Surveillance is essential for detecting outbreaks early, identifying high-risk neighborhoods, and guiding vaccination or WASH interventions.


Figure 6.0

Figure 6.0
Clinical Nursing Flow Sheet for Suspected Typhoid Case Management



Figure 7.0


Community Typhoid Outbreak Response Flow Sheet



Note. This flow sheet summarizes the community response pathway for typhoid outbreak control.

Figure 8.0
Nursing Role Flow Sheet in Typhoid Outbreak Control

Nursing Role in Typhoid Outbreak Response



Note. This flow sheet summarizes the integrated nursing roles in typhoid outbreak control.

Table 2.0

Nursing Roles in Typhoid Outbreak Response

Nursing role	Main activities	Expected outcome
Clinical care	Triage, vital signs, hydration, fever monitoring, medication administration	Early recognition and improved recovery
Infection prevention and control	Hand hygiene, safe disposal of stool/vomit, surface cleaning, PPE use	Reduced transmission in healthcare settings
Vaccination support	TCV administration, cold-chain monitoring, documentation, adverse event monitoring	Improved vaccine coverage
Health education	Safe water, food hygiene, handwashing, antibiotic adherence	Improved prevention practices
Community mobilization	Door-to-door visits, school awareness, family counseling	Increased vaccine acceptance and early care-seeking
Surveillance	Fever case identification, data recording, public health reporting	Early outbreak detection and response
Preparedness	Training, outbreak team participation, emergency planning	Stronger outbreak readiness

Table 3.0

Major Barriers to Typhoid Control in Karachi and Nursing Responses

Barrier	Contribution to typhoid transmission	Nursing response
Unsafe drinking water	Increases fecal–oral spread	Teach boiling, chlorination, and safe storage
Poor sanitation	Contaminates household and community environments	Promote latrine use and safe waste disposal
Street-vended unsafe food	Increases exposure to contaminated food	Counsel families and vendors on safe food handling
Overcrowded settlements	Facilitates rapid spread	Conduct household screening and education
Antibiotic misuse	Promotes resistance and treatment failure	Educate on prescribed antibiotics and adherence
Vaccine hesitancy	Reduces campaign coverage	Address myths and counsel families
Weak surveillance	Delays outbreak detection	Report fever clusters and confirmed cases
Low awareness	Delays care-seeking	Conduct school and community sessions

Table 4.0

Evidence-to-Practice Implications for Nursing

Evidence area	Nursing implication	Practice recommendation
XDR typhoid burden	Nurses must recognize resistant typhoid risk	Strengthen triage and early reporting
High pediatric burden	Children require priority prevention	Support school-based and community vaccination
Unsafe WASH conditions	Transmission continues without hygiene change	Integrate WASH education into every contact
Vaccine effectiveness	TCV reduces typhoid risk	Maintain high TCV coverage and mop-up campaigns
Antibiotic resistance	Misuse worsens XDR spread	Counsel against self-medication
Karachi outbreak concentration	Urban slums need targeted action	Deploy nurse-led outreach in high-risk communities

Discussion

The findings show that typhoid control in Karachi requires an integrated response involving clinical care, vaccination, surveillance, antimicrobial stewardship, and WASH interventions. The XDR outbreak demonstrated that antimicrobial resistance can rapidly transform a treatable infection into a major public health emergency. Karachi's role as a major outbreak setting highlights the vulnerability of densely populated cities with inadequate sanitation and unsafe water systems.

Nurses were central to the response at multiple levels. In clinical settings, nurses identified suspected cases, monitored patients, administered treatment, prevented dehydration, supported infection control, and educated families. In vaccination campaigns, nurses and Lady Health Workers helped operationalize TCV delivery, manage records, support cold-chain processes, monitor adverse events, and build community acceptance. In communities, nurses promoted safe water, handwashing, food hygiene, and timely care-seeking.

A key lesson is that outbreak response cannot rely only on physicians, laboratories, or vaccines. Nurses are the bridge between health systems and communities. Their trusted presence allows them to communicate prevention messages, identify early cases, reduce vaccine hesitancy, and reinforce safe practices. However, nursing roles must be formally recognized, documented, and evaluated in outbreak response plans.

The review also highlights the need for stronger public health nursing capacity in Pakistan. Nurses should be trained in outbreak investigation, risk communication, infection prevention, surveillance, vaccine counseling, and antimicrobial stewardship. They should be included in rapid response teams and epidemic task forces at hospital, district, and provincial levels.

Recommendations

1. Strengthen public health nursing roles in typhoid surveillance, vaccination, and community education.
2. Train nurses in XDR typhoid management, including early recognition, IPC, and antimicrobial stewardship.

3. Integrate nurses into outbreak response teams at hospital and district levels.
4. Expand nurse-led school and community education on handwashing, safe water, food hygiene, and early care-seeking.
5. Maintain high TCV coverage through routine immunization and targeted mop-up campaigns.
6. Improve WASH infrastructure in high-risk Karachi settlements.
7. Develop nursing documentation tools for typhoid case reporting, vaccination follow-up, and household education.
8. Conduct research on nursing outcomes in typhoid outbreak response to quantify the impact of nursing-led interventions.

Limitations

This review used a narrative synthesis because included sources varied in design, quality, and focus. Some nursing contributions were described in outbreak reports and grey literature rather than formal research studies. There is limited quantitative evidence measuring the direct impact of nurse-led interventions on typhoid incidence. Some case trend values used for graphs are approximate and should be verified against final datasets before journal submission.

Conclusion

Typhoid fever, particularly XDR typhoid, remains a serious public health challenge in Karachi and Pakistan. The Sindh outbreak demonstrated how unsafe water, poor sanitation, overcrowding, antimicrobial resistance, and delayed care can combine to create a sustained epidemic. Nurses have played a critical role in controlling typhoid through clinical care, infection prevention, vaccination, surveillance, health education, and community mobilization.

To reduce typhoid transmission and improve outbreak preparedness, Pakistan must strengthen nursing capacity, sustain TCV coverage, improve WASH conditions, promote rational antibiotic use, and integrate nurses into public health decision-making. A nurse-enabled outbreak response model can help Karachi and other endemic regions move toward sustainable typhoid control.

REFERENCES

- Appiah, G. D., Chung, A., Bentsi-Enchill, A. D., Kim, S., Crump, J. A., Mogasale, V., Pellegrino, R., Slayton, R. B., & Mintz, E. D. (2020). Typhoid outbreaks, 1989–2018: Implications for prevention and control. *The American Journal of Tropical Medicine and Hygiene*, 102(6), 1296–1305. <https://doi.org/10.4269/ajtmh.19-0624>
- Aslam, F., Yue, Y., & Aziz, M. (2021). Introduction of typhoid vaccine in the expanded immunization program of Pakistan. *Human Vaccines & Immunotherapeutics*, 17(7), 2132. <https://doi.org/10.1080/21645515.2020.1869496>
- Centers for Disease Control and Prevention. (2023). *Typhoid and paratyphoid fever*. CDC Yellow Book.
- Dyson, Z. A., Klemm, E. J., Palmer, S., & Dougan, G. (2019). Antibiotic resistance and typhoid. *Clinical Infectious Diseases*, 68(Suppl. 2), S165–S170. <https://doi.org/10.1093/cid/ciy1111>
- Fatima, M., Kumar, S., Hussain, M., Memon, N. M., Vighio, A., Syed, M. A., Chaudhry, A., Hussain, Z., Baig, Z. I., Baig, M. A., Asghar, R. J., Ikram, A., & Khader, Y. (2021). Morbidity and mortality associated with typhoid fever among hospitalized patients in Hyderabad District, Pakistan, 2017–2018: Retrospective record review. *JMIR Public Health and Surveillance*, 7(5), Article e27268. <https://doi.org/10.2196/27268>

- GBD 2017 Typhoid and Paratyphoid Collaborators. (2019). The global burden of typhoid and paratyphoid fevers: A systematic analysis for the Global Burden of Disease Study 2017. *The Lancet Infectious Diseases*, 19(4), 369–381. [https://doi.org/10.1016/S1473-3099\(18\)30685-6](https://doi.org/10.1016/S1473-3099(18)30685-6)
- Gavi, the Vaccine Alliance. (2023). *How a major typhoid vaccine campaign helped Nepal find missed-out “zero-dose” kids*. Gavi.
- Hancuh, M., Walldorf, J., Minta, A. A., Tevi-Benissan, C., Christian, K. A., Nedelec, Y., Heitzinger, K., Mikoleit, M., Tiffany, A., Bentsi-Enchill, A. D., & Breakwell, L. (2023). Typhoid fever surveillance, incidence estimates, and progress toward typhoid conjugate vaccine introduction—Worldwide, 2018–2022. *Morbidity and Mortality Weekly Report*, 72(7), 171–176. <https://doi.org/10.15585/mmwr.mm7207a2>
- John, J., Bavdekar, A., Rongsen-Chandola, T., Dutta, S., Kang, G., & the NSSEFI Collaborators. (2023). Burden of typhoid and paratyphoid fever in India. *The New England Journal of Medicine*, 388(16), 1491–1500. <https://doi.org/10.1056/NEJMoa2209449>
- Khan, M. I., Soofi, S. B., Ochiai, R. L., Khan, M. J., Sahito, S. M., Habib, M. A., Puri, M. K., von Seidlein, L., Clemens, J. D., & Bhutta, Z. A. (2015). Typhoid vaccine introduction: An evidence-based pilot implementation project in Nepal and Pakistan. *Vaccine*, 33(Suppl. 3), C62–C67. <https://doi.org/10.1016/j.vaccine.2015.03.080>
- Khawaja, S. (2019). *Prevention in action: Protecting communities against typhoid in Pakistan*. Coalition Against Typhoid.
- Kuehn, R., Stoesser, N., Eyre, D., Darton, T. C., & Basnyat, B. (2025). Enteric fever. *The Lancet*.
- Mogasale, V., Maskery, B., Ochiai, R. L., Lee, J. S., Mogasale, V. V., Ramani, E., Kim, Y. E., Park, J. K., & Wierzb, T. F. (2014). Burden of typhoid fever in low-income and middle-income countries: A systematic, literature-based update with risk-factor adjustment. *The Lancet Global Health*, 2(10), e570–e580. [https://doi.org/10.1016/S2214-109X\(14\)70301-8](https://doi.org/10.1016/S2214-109X(14)70301-8)
- National Institute of Health Pakistan. (2022). *Field epidemiology and disease surveillance reports on enteric fever and antimicrobial resistance*. Government of Pakistan.
- Pradhan, N. A., & Gavi, the Vaccine Alliance. (2023). *Nepal’s typhoid conjugate vaccine campaign: A catalyst to reach zero-dose children*. Coalition Against Typhoid.
- Qamar, F. N., Yousafzai, M. T., Sultana, S., Baig, A., Memon, H., Junejo, A., Kazi, A. M., & Garrett, D. O. (2018). A retrospective study of laboratory-based enteric fever surveillance, Pakistan, 2012–2014. *The Journal of Infectious Diseases*, 218(Suppl. 4), S201–S205. <https://doi.org/10.1093/infdis/jiy205>
- Rane, S. (2011). Street vended food in developing world: Hazard analyses. *Indian Journal of Microbiology*, 51(1), 100–106. <https://doi.org/10.1007/s12088-011-0154-x>
- Stanaway, J. D., Reiner, R. C., Blacker, B. F., Goldberg, E. M., Khalil, I. A., Troeger, C. E., Andrews, J. R., Bhutta, Z. A., Crump, J. A., Im, J., Marks, F., Mintz, E., Park, S. E., Zaidi, A. K. M., Abebe, Z., Abejie, A. N., Adedje, I. A., Ali, B. A., Amare, A. T., ... Mokdad, A. H. (2019). The global burden of typhoid and paratyphoid fevers: A systematic analysis for the Global Burden of Disease Study 2017. *The Lancet Infectious Diseases*, 19(4), 369–381. [https://doi.org/10.1016/S1473-3099\(18\)30685-6](https://doi.org/10.1016/S1473-3099(18)30685-6)
- Tharwani, Z. H., Kumar, P., Salman, Y., Islam, Z., Ahmad, S., & Essar, M. Y. (2022). Typhoid in Pakistan: Challenges, efforts, and recommendations. *Infection and Drug Resistance*, 15, 2523–2527. <https://doi.org/10.2147/IDR.S365220>

- UNICEF Nepal. (2022). *Nepal introduces typhoid vaccine into routine immunisation across the country*. UNICEF.
- World Health Organization. (2018). *Typhoid vaccines: WHO position paper—March 2018*. *Weekly Epidemiological Record*, 93(13), 153–172.
- World Health Organization. (2018). *Typhoid fever—Islamic Republic of Pakistan*. Disease Outbreak News.
- World Health Organization. (2023). *Typhoid*. World Health Organization.
- World Health Organization. (2024). *State of the world's nursing and health workforce priorities*. World Health Organization.
- Yousafzai, M. T., Karim, S., Qureshi, S., Kazi, M., Memon, H., Junejo, A., & Khawaja, Z. (2021). Effectiveness of typhoid conjugate vaccine against culture-confirmed *Salmonella enterica* serotype Typhi in an extensively drug-resistant outbreak setting of Hyderabad, Pakistan: A cohort study. *The Lancet Global Health*, 9(8), e1154–e1162. [https://doi.org/10.1016/S2214-109X\(21\)00255-2](https://doi.org/10.1016/S2214-109X(21)00255-2)