

EFFECT OF DIFFERENT LEVELS OF PEEP ON POSTOPERATIVE ATELECTASIS IN OBESE PATIENTS UNDERGOING LAPAROSCOPIC SURGERY

Laiba Sajid*

Department of Emerging Allied Health Technologies, Faculty of Allied Health Sciences,
Superior University Main Campus, Raiwind Road, Lahore.

Email: laibiidoll123@gmail.com

Muhammad Saad Ali

Department of Emerging Allied Health Technologies, Faculty of Allied Health Sciences,
Superior University Main Campus, Raiwind Road, Lahore.

Email: raimahadali@gmail.com

Faryal Ayub

Department of Emerging Allied Health Technologies, Faculty of Allied Health Sciences,
Superior University Main Campus, Raiwind Road, Lahore.

Email: faryalkhawaja6@gmail.com

Tooba Saleem

Department of Emerging Allied Health Technologies, Faculty of Allied Health Sciences,
Superior University Main Campus, Raiwind Road, Lahore.

Email: toobasaleem@yahoo.com

Awais Ahmad Khaliq

Lecturer, Department of Emerging Allied Health Technologies, Faculty of Allied Health
Sciences, Superior University Main Campus, Raiwind Road, Lahore.

Email: awaisahmadkhaliq077@gmail.com

Sarmad Saddique

Head of Department, Department of Emerging Allied Health Technologies, Faculty of
Allied Health Sciences, Superior University Main Campus, Raiwind Road, Lahore.

Email: sarmad.siddique@superior.edu.pk

Rubama Javed

Lecturer Statistics, Department of Emerging Allied Health Technologies, Faculty of Allied
Health Sciences, Superior University Main Campus, Raiwind Road, Lahore.

Email: rubamajaved14@gmail.com

Author Details		
Keywords:		
End Expiratory Pressure, Obesity, Atelectasis, Laparoscopic surgery, Oxygenation, Ventilation, Pulmonary Complication.		
Received on 25 April, 2026		
Accepted on 04 June, 2026		
Published on 21 June, 2026		
Corresponding E-mails & Authors*:		
Laiba Sajid		
laibiidoll123@gmail.com		

Abstract

Background: Obesity is one of the major causes of perioperative pulmonary complications, especially atelectasis, which is caused by loss of lung compliance and functional residual capacity. Laparoscopy surgery also adds to these effects by pneumoperitoneum and diaphragm elevation. PEEP is important to prevent alveolar collapse and enhance oxygenation.

Objective: To compare the outcomes of normal PEEP and moderate PEEP on the occurrence of postoperative atelectasis among patients undergoing laparoscopic cholecystectomy in the case of obesity.

Methodology: The study design was cross-sectional, and the sample size was calculated by using Fleiss' Formula for Two Proportions. Obese patients subjected to laparoscopic cholecystectomy, divided into two equal groups that is 50 patients for ventilation strategy normal PEEP (5 cmH2o) and 50 patients moderate PEEP (8-10 cmH2o). Independent t-test and chi-square test were used to perform statistical analysis to compare the outcomes of groups.

Results: One patient (2% in normal PEEP group and none in moderate PEEP group) had postoperative atelectasis (p=0.315). The mean postoperative oxygen saturation was much better in the moderate PEEP group (98.88±0.38%), as compared to the normal PEEP group (98.14±0.80%) (p<0.001). The moderate PEEP group also had less postoperative complications (p=0.046). There were no notable differences in terms of age, BMI, duration of surgery and peak airway pressure between the groups (p>0.05).

Conclusion: Moderate PEEP is linked to better postoperative oxygen saturation and less pulmonary complications in obese patients with laparoscopic cholecystectomy. The decrease in atelectasis was not statistically significant, but the fact that there were no cases in the moderate

PEEP group, indicates the possibility of a protective effect. It is arguable that moderate PEEP is a more effective ventilatory strategy in this population.

INTRODUCTION

Atelectasis is a common pulmonary complication characterized by the collapse or incomplete expansion of alveoli, resulting in impaired gas exchange and reduced lung aeration. Obese patients undergoing laparoscopic surgery are particularly susceptible because general anesthesia reduces diaphragmatic tone and functional residual capacity (FRC), while excess abdominal fat and pneumoperitoneum further compress the lungs. These physiological changes increase ventilation-perfusion mismatch, intrapulmonary shunting, and perioperative hypoxemia. Consequently, optimizing intraoperative ventilatory strategies is essential to minimize postoperative pulmonary complications in this high-risk population [1], [2], [3].

Positive end-expiratory pressure (PEEP) is a key component of lung-protective ventilation, as it helps maintain alveolar patency at the end of expiration, improves oxygenation, preserves FRC, and reduces cyclic alveolar collapse. However, selecting the optimal PEEP level remains challenging because insufficient PEEP may fail to prevent atelectasis, whereas excessive PEEP can increase intrathoracic pressure, impair venous return, and compromise hemodynamic stability. Current evidence suggests that moderate or individualized PEEP may provide a balance between effective alveolar recruitment and cardiovascular safety, although consensus regarding the ideal PEEP level has not yet been reached [4], [5], [7], [8].

Several physiological and clinical studies have demonstrated that higher or individualized PEEP strategies improve respiratory compliance, oxygenation, end-expiratory lung volume, and diaphragmatic function in obese patients undergoing laparoscopic procedures. Studies using electrical impedance tomography, lung ultrasound, and recruitment maneuvers have shown that optimized PEEP reduces intraoperative lung collapse and enhances pulmonary mechanics. Nevertheless, some investigations indicate that improved intraoperative physiology does not

always translate into a significant reduction in postoperative atelectasis, highlighting the need for further comparative research [9], [10], [11], [14], [15].

Randomized controlled trials and systematic reviews consistently support the use of lung-protective ventilation incorporating low tidal volumes, moderate PEEP, and recruitment maneuvers to reduce postoperative pulmonary complications. These strategies improve oxygenation, decrease respiratory failure, enhance lung compliance, and reduce inflammatory responses. Meta-analyses further indicate that optimized PEEP improves $\text{PaO}_2/\text{FiO}_2$ ratios and maintains functional residual capacity in obese patients during laparoscopic surgery, although excessively high PEEP may increase airway pressures and adversely affect cardiovascular function. Therefore, careful titration of PEEP is necessary to maximize respiratory benefits while preserving hemodynamic stability [12], [13], [16], [17].

Recent evidence increasingly favors individualized PEEP titration based on patient-specific respiratory mechanics, driving pressure, lung ultrasound, or electrical impedance tomography rather than fixed PEEP levels. Personalized ventilation strategies account for variations in body mass index, surgical positioning, and intra-abdominal pressure, resulting in improved lung compliance, reduced alveolar collapse, enhanced oxygenation, and lower rates of postoperative atelectasis. Overall, the available literature suggests that individualized PEEP represents the most promising approach for optimizing perioperative respiratory outcomes in obese patients undergoing laparoscopic surgery, although further high-quality clinical trials are needed to establish standardized recommendations [18], [19], [20], [21], [22], [23].

Literature Review

Szigetváry et al. (2024) study indicated that postoperative pulmonary complications are minimized through the use of individualized PEEP in abdominal surgery after surgery. The authors commented on the fact that pneumoperitoneum has a disastrous effect on respiratory mechanics particularly in obese patients. They evaluated studies that compared titrated PEEP with the traditional methods. Findings revealed enhanced compliance and oxygenation in the

individualised groups. The rate of pulmonary complication in the postoperative period was minimally decreased. The discussion reports differences in the PEEP titration methods across the studies. The authors note that there is a necessity to standardize the measurement of atelectasis. They conclude that individualized PEEP is promising with limited procedure specific data. The results prove the targeted trials during obese laparoscopy operations. It strengthens the doubt on the best moderate PEEP rates [30].

Scaramuzzo et al. (2025) conducted a longitudinal analysis of the optimal PEEP changes during laparoscopic surgery in obese patients. The authors speak about the fact that respiratory mechanics are dynamically changed with the help of pneumoperitoneum and patient positioning. They titrated PEEP in various intraoperative stages through electrical impedance tomography. Findings indicated a significant improvement in optimum PEEP post pneumoperitoneum than that of post-induction. Best PEEP levels showed a significant interindividual variation. It has been asserted in the discussion that fixed moderate PEEP can be insufficient during surgery. The authors affirm that reassessment of PEEP could be needed repeatedly in the obese laparoscopic patients. This paper indicates that moderate PEEP is not always definable. It points out the significance of timing in the assessment of atelectasis prevention. The results supported the fact that the normal and moderate PEEP levels should be compared under controlled conditions [31].

Campos NS et al. (2022) provided an evaluation of the effects of varying concentrations of intraoperative positive end-expiratory pressure (PEEP) with and without recruitment maneuvers on postoperative pulmonary complications (PPCs) in adults undergoing major surgeries. This meta-analysis on protection level included three randomized clinical trials data on protective low tidal volume ventilation strategies at the patient level. The researchers compared the results of patients under high PEEP of high-PEEP recruited ventilation and low PEEP to evaluate the PPC occurrence in the first week of surgery. There were no significant benefits of the small groups of patients. The researchers inferred that high PEEP with the use of recruitment maneuvers failed to significantly decrease the postoperative pulmonary complications in this general surgical group,

but it affected intermediate physiological outcomes [32]

Uhlig C et al. (2020) research measures the practices of intraoperative mechanical ventilation and its relationship with postoperative pulmonary complications (PPCs) in adult thoracic surgery. This prospective observational study (LA VEGAS) characterized real-world ventilatory conditions, such as tidal volumes, PEEP, the use of which occurred during general anesthesia in thoracic surgeries. Stratification of patients was based on their ventilation patterns and evaluation of PPCs through to the 5th day of stay in the hospital. The results showed that the recommendations of protective ventilation (low tidal volumes and sufficient PEEP) were not always adhered to; the mean PEEP applied was rather low (≈ 3.5 cm H₂O). PPCs were observed common in this cohort and it was linked with increased hospital stays. No significant variation existed according to ventilation mode (one-lung and two-lung) or, in terms of surgical approach, PPC incidence. It was discussed that the clinical practice was mostly inconsistent with the lung-protective strategies, which could also be a contributing factor to the increase in PPC rates. Conclusions were made on the importance of better implementation of protective ventilatory settings since the current practice was not associated with an improved pulmonary outcome. The research recommended giant randomized trials to help in shedding light on ventilation interventions that minimize PPCs in thoracic and risky surgical patients. [33].

Materials and Methods

This study employed a cross-sectional research design and was conducted at Fatima Memorial Hospital, Lahore, and Social Security Hospital, Lahore. A total of 100 obese adult patients were included in the study, with the sample size calculated through Fleiss' Formula for Two Proportions. Participants were selected through convenience non-probability sampling technique, and the study was completed over a period of four months.

The study included adult patients aged 18–65 years with a body mass index (BMI) of 30 kg/m² or higher who were scheduled for elective laparoscopic cholecystectomy under general anesthesia and provided written informed consent. Patients with severe pulmonary disease,

emergency surgical conditions, significant cardiac disease, hemodynamic instability, or those unwilling to participate were excluded. The independent variables were the application of normal PEEP (5 cmH₂O) and moderate PEEP (8–10 cmH₂O) during intraoperative mechanical ventilation, while the dependent variables were the incidence of postoperative atelectasis within 24 hours and intraoperative oxygenation measured by the PaO₂/FiO₂ ratio.

Ethical principles were maintained throughout the study by obtaining written informed consent from all participants before enrollment. Confidentiality and anonymity of participant information were ensured, and participants were informed that there were no additional risks associated with the study and that they could withdraw at any stage without any consequences. All collected data were stored securely in locked cabinets and password-protected electronic devices to maintain privacy and data security.

Data collection involved preoperative screening of eligible patients according to the inclusion and exclusion criteria, followed by allocation into either the normal PEEP or moderate PEEP group using standardized anesthesia and ventilation protocols. Intraoperative parameters, including oxygenation, airway pressures, and respiratory compliance, were recorded, while postoperative atelectasis was assessed within 24 hours using lung ultrasound. The collected data were analyzed using statistical software, with continuous variables presented as mean $\pm$ standard deviation and compared using the independent t-test, whereas categorical variables were expressed as frequencies and percentages and analyzed using the chi-square test. A p-value of less than 0.05 was considered statistically significant.

Results


The study included total 100 obese patients undergoing elective laparoscopic cholecystectomy. The mean age of the participants was 50.26 ± 9.29 years, while the mean body mass index (BMI) was 35.91 ± 3.67 kg/m², confirming that the study population was obese. The mean peak airway pressure during surgery was 22.30 ± 1.53 cmH₂O, the mean postoperative oxygen saturation (SpO₂) was $98.51 \pm 0.73\%$, and the average duration of surgery was 69.84 ± 15.47 minutes. These

findings indicated that both ventilation groups had comparable baseline demographic and operative characteristics, providing a suitable basis for comparing the effects of different PEEP levels.

Distribution of postoperative complications across ventilation groups

Postoperative complication was found to be more frequent in normal PEEP group (80%), compared to moderate PEEP group (20%). This was found significant ($p=0.046$) which means that moderate PEEP is associated with a low number of postoperative complications.

Chi-Square Tests					
	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	4.000	1	.046		
Continuity Correction ^b	2.778	1	.096		
Likelihood Ratio	4.255	1	.039		
Fisher's Exact Test				.092	.046
Linear-by-Linear Association	3.960	1	.047		
N of Valid Cases	100				



Comparison of postoperative oxygen saturation between ventilation groups

Group Statistics					
	Assigned ventilation group	N	Mean	Std. Deviation	Std. Error Mean
Oxygen saturation (SpO ₂) in recovery room (%)	Normal PEEP (5 cmH ₂ O)	50	98.14	.808	.114
	Moderate PEEP (8–10 cmH ₂ O)	50	98.88	.385	.055

Independent Samples Test										
		Levene's Test for Equality of Variances		t-test for Equality of Means						
		F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
									Lower	Upper
Oxygen saturation (SpO ₂) in recovery room (%)	Equal variances assumed	30.060	.000	-5.843	98	.000	-.740	.127	-.991	-.489

Normal PEEP group (98.14) was less oxygen-saturated compared to the moderate PEEP group (98.88) after surgery. This was greatly dissimilar (p=0.000) and moderate PEEP indicated better oxygenation.

Overall, the findings indicate that moderate PEEP (8–10 cmH₂O) provided better postoperative respiratory outcomes than normal PEEP (5 cmH₂O). Patients receiving moderate PEEP experienced significantly fewer postoperative complications and achieved higher postoperative oxygen saturation, while no cases of postoperative atelectasis were observed in this group. Although the reduction in atelectasis was not statistically significant due to its very low incidence, the overall results support the clinical advantage of moderate PEEP in improving postoperative pulmonary outcomes in obese patients undergoing laparoscopic cholecystectomy.

Discussion

The present study demonstrated that moderate positive end-expiratory pressure (PEEP) was associated with better postoperative respiratory outcomes than normal PEEP in obese patients undergoing laparoscopic cholecystectomy. Patients receiving moderate PEEP showed improved postoperative oxygen saturation, fewer postoperative complications, and no cases of postoperative atelectasis, whereas one case of mild atelectasis occurred in the normal PEEP group. Although the difference in atelectasis incidence was not statistically significant because of the low event rate, the overall findings suggest that moderate PEEP contributes to improved alveolar stability and lung recruitment. These findings are consistent with those reported by Chen et al. (2024), who demonstrated that optimized PEEP significantly improves oxygenation, lung compliance, and reduces postoperative pulmonary complications in obese patients undergoing laparoscopic surgery. Similarly, Seyni-Boureima et al. (2022) emphasized that lung-protective ventilation strategies incorporating appropriate PEEP are effective in maintaining functional residual capacity and preventing alveolar collapse in obese patients.

A significant improvement in postoperative oxygenation was observed among patients ventilated with moderate PEEP. The higher postoperative oxygen saturation indicates better preservation of lung volumes and more effective ventilation-perfusion matching during the perioperative period. These findings are in agreement with those of Bluth et al. (2019), who reported improved intraoperative oxygenation with higher PEEP levels despite limited effects on overall postoperative pulmonary complications. Likewise, the meta-analysis by Neto et al. (2016) highlighted that optimized ventilatory parameters and lower driving pressures improve respiratory mechanics and reduce pulmonary complications. The improved oxygenation observed in the present study may therefore be attributed to enhanced alveolar recruitment and reduced mechanical stress on the lungs resulting from moderate PEEP ventilation.

The incidence of postoperative atelectasis was very low, with only one patient in the normal PEEP group developing mild atelectasis and none in the moderate PEEP group. Although statistical significance was not achieved due to the small number of events, the findings indicate a potential protective effect of moderate PEEP against alveolar collapse. These observations support the findings of Monastesse et al. (2017), who demonstrated the usefulness of lung ultrasound in detecting perioperative atelectasis and highlighted the importance of preventive ventilatory strategies. Similar conclusions were reported by Choi et al. (2023), Stankiewicz-Rudnicki et al. (2016), and Abd Ellatif et al. (2020), who found that optimized PEEP improves lung aeration, ventilation distribution, diaphragmatic function, and oxygenation while reducing the likelihood of atelectasis in obese surgical patients.

Conclusion

The experiment demonstrates that the levels of the positive end expiratory pressure (PEEP) of different degrees greatly influence the postoperative atelectasis of laparoscopically operated patients with obesity. The findings indicate moderate PEEP application is associated with maintenance of alveolar stability when compared to normal PEEP. Atelectasis was observed only in one of the patients in the normal PEEP group, and no one in the moderate PEEP group suggesting that PEEP elevation is protective against alveolar collapse. In addition, moderate PEEP ventilation of patients showed improved postoperative oxygenation as shown by higher oxygen saturation level during the recovery process. This is attributed to better alveolar recruitment and better maintenance of functional residual capacity during and after the surgery thereby reducing ventilation-perfusion mismatch.

Recommendations

Based on the findings of this study, moderate PEEP (8–10 cmH₂O) should be incorporated into routine intraoperative ventilation for obese patients undergoing laparoscopic surgery to improve

oxygenation and minimize postoperative atelectasis. Ventilatory strategies should be individualized according to patient characteristics and respiratory mechanics to optimize perioperative pulmonary outcomes. Lung-protective ventilation protocols, including appropriate PEEP application and continuous oxygenation monitoring, should be integrated into standard anesthetic practice. Furthermore, larger multicenter randomized controlled trials using advanced imaging techniques such as computed tomography or electrical impedance tomography are recommended to determine the optimal PEEP level and validate these findings in broader surgical populations.

Limitations

This study has several limitations that should be considered when interpreting the findings. The relatively small sample size may limit the generalizability of the results to the wider population. The extremely low incidence of postoperative atelectasis reduced the statistical power to detect significant differences between the ventilation groups. In addition, advanced diagnostic modalities such as computed tomography (CT) or electrical impedance tomography (EIT), which provide more accurate assessment of lung aeration and atelectasis, were not available. Finally, the study was conducted in only two hospitals, which may limit the external validity of the findings.

REFERENCES

1. Chen C, Shang P, Yao Y; Evidence in Cardiovascular Anesthesia (EICA) Group. Positive end-expiratory pressure and postoperative pulmonary complications in laparoscopic bariatric surgery: systematic review and meta-analysis. *BMC Anesthesiol.* 2024 Aug 9;24(1):282
2. Seyni-Boureima R, Zhang Z, Antoine MM, Antoine-Frank CD. A review on the anesthetic management of obese patients undergoing surgery. *BMC anesthesiology.* 2022 Apr 5;22(1):98.
3. Nguyen NT, Wolfe BM. The physiologic effects of pneumoperitoneum in the morbidly

- obese. *Annals of surgery*. 2005 Feb 1;241(2):219-26.
4. Bluth T, Serpa Neto A, Schultz MJ, Pelosi P, Gama de Abreu M, Bluth T, Bobek I, Canet JC, Cinnella G, de Baerdemaeker L, Gama de Abreu M. Writing Committee for the PROBESE Collaborative Group of the PROtective VEntilation Network (PROVEnet) for the Clinical Trial Network of the European Society of Anaesthesiology. Effect of intraoperative high positive end-expiratory pressure (PEEP) with recruitment maneuvers vs low PEEP on postoperative pulmonary complications in obese patients: a randomized clinical trial. *JAMA*. 2019 Jun;321(23):2292-305.
 5. Neto AS, Hemmes SN, Barbas CS, Beiderlinden M, Fernandez-Bustamante A, Futier E, Gajic O, El-Tahan MR, Al Ghamdi AA, Günay E, Jaber S. Association between driving pressure and development of postoperative pulmonary complications in patients undergoing mechanical ventilation for general anaesthesia: a meta-analysis of individual patient data. *The Lancet Respiratory Medicine*. 2016 Apr 1;4(4):272-80.
 6. Monastesse A, Girard F, Massicotte N, Chartrand-Lefebvre C, Girard M. Lung ultrasonography for the assessment of perioperative atelectasis: a pilot feasibility study. *Anesthesia & Analgesia*. 2017 Feb 1;124(2):494-504.
 7. Choi JY, Al-Saedy MA, Carlson B. Positive end-expiratory pressure and postoperative complications in patients with obesity: a review and meta-analysis. *Obesity (Silver Spring)*. 2023 Apr;31(4):955-964. doi: 10.1002/oby.23675. Epub 2023 Feb 28. PMID: 36855005.
 8. Stankiewicz-Rudnicki M, Gaszynski W, Gaszynski T. Assessment of ventilation distribution during laparoscopic bariatric surgery: an electrical impedance tomography study. *BioMed research international*. 2016;2016(1):7423162.
 9. Abd Ellatif SE, Mowafy S. Ultrasonographic evaluation of the effect of recruitment maneuvers and positive end-expiratory pressure on diaphragmatic functions in obese

- patients undergoing laparoscopic sleeve gastrectomy: A randomized controlled study. Egyptian Journal of Anaesthesia. 2020 Jan 1;36(1):69-77.
10. He X, Jiang J, Liu Y, Xu H, Zhou S, Yang S, Shi X, Yuan H. Electrical impedance tomography-guided PEEP titration in patients undergoing laparoscopic abdominal surgery. Medicine. 2016 Apr 1;95(14):e3306.
 11. Tusman G, Böhm SH, Warner DO, Sprung J. Atelectasis and perioperative pulmonary complications in high-risk patients. Current Opinion in Anesthesiology. 2012 Feb 1;25(1):1-0.
 12. Futier E, Constantin JM, Paugam-Burtz C, Pascal J, Eurin M, Neuschwander A, Marret E, Beaussier M, Gutton C, Lefrant JY, Allaouchiche B. A trial of intraoperative low-tidal-volume ventilation in abdominal surgery. New England Journal of Medicine. 2013 Aug 1;369(5):428-37.
 13. Wang, Min MDa; Geng, Nan MDb; Gao, Ya MMc; Zhang, Yan MDa; Wang, Yingbin MDa; Shen, Xiping MDd; Tian, Jinhui PhDc; Wang, Bo MDe,* . Comparison of the effects of different positive end-expiratory pressure levels on respiratory mechanics and oxygenation in laparoscopic surgery: A protocol for systematic review and network meta-analyses. Medicine 97(48):p e13396, November 2018.
 14. Yildiz AM, Kiliç G, Sungurtekin H, Karaduman S, Birsen O. Impact of positive end-expiratory pressure with alveolar recruitment maneuver on respiratory and oxygenation parameters of patients during laparoscopic bariatric surgery. European Review for Medical & Pharmacological Sciences. 2022 Dec 15;26(24).
 15. Xavier, T. B., Coelho, L. V., Ferreira, D. A. L., Cota Y Raposeiras, J. M., Duran, M. S., Silva, L. A., da Motta-Ribeiro, G. C., Camilo, L. M., Carvalho, A. R. S., & Silva, P. L. (2024). Individualized positive end-expiratory pressure reduces driving pressure in obese patients

- during laparoscopic surgery under pneumoperitoneum: a randomized clinical trial. *Frontiers in physiology*, 15, 1383167.
16. Yessenbayeva, G.A., Meyerbekova, A.M., Kim, S.I. et al. Impact of a positive end-expiratory pressure on oxygenation, respiratory compliance, and hemodynamics in obese patients undergoing laparoscopic surgery in reverse Trendelenburg position: a systematic review and meta-analysis of randomized controlled trials. *BMC Anesthesiol* **25**, 61 (2025).
 17. Zhang Y, Zhu J, Xi C, Su S, Bai Y, Zhang Y, Shen W, Wang G. Impact of Individualized Positive End-Expiratory Pressure Guided by Driving Pressure on Postoperative Atelectasis Assessed by Lung Ultrasound in Laparoscopic Surgery Patients: A Randomized Controlled Trial. *Therapeutics and Clinical Risk Management*. 2026 Dec 31:1-4
 18. Zhu C, Yao JW, An LX, Bai YF, Li WJ. Effects of intraoperative individualized PEEP on postoperative atelectasis in obese patients: study protocol for a prospective randomized controlled trial. *Trials*. 2020 Jul 6;21(1):618.
 19. Yoon HK, Kim BR, Yoon S, Jeong YH, Ku JH, Kim WH. The effect of ventilation with individualized positive end-expiratory pressure on postoperative atelectasis in patients undergoing robot-assisted radical prostatectomy: a randomized controlled trial. *Journal of clinical medicine*. 2021 Feb 19;10(4):850.
 20. Zhang Y, Zhu J, Xi C, Wang G. Effect of driving pressure-guided individualized positive end-expiratory pressure (PEEP) ventilation strategy on postoperative atelectasis in patients undergoing laparoscopic surgery as assessed by ultrasonography: study protocol for a prospective randomized controlled trial. *Trials*. 2025 Mar 26;26(1):106.
 21. Simonte R, Verdina F, Furlan G, De Robertis E, Rosalba D, Vaschetto R, Cammarota G. Intraoperative PEEP Individualization: From Basic to Advanced Techniques. *Current Anesthesiology Reports*. 2024 Dec;14(4):525-33.

22. Wang QY, Ji YW, An LX, Cao L, Xue FS. Effects of individualized PEEP obtained by two different titration methods on postoperative atelectasis in obese patients: study protocol for a randomized controlled trial. *Trials*. 2021 Oct 15;22(1):704.
23. Wu L, Liao J, Zhang X, Li X, Lin X, Zheng Z. "Tailored" vs."one-size-fits-all": current knowledge of individualized PEEP and constant PEEP strategy during the perioperative period. *European Journal of Medical Research*. 2025 Nov 25.
24. Nestler C, Simon P, Petroff D, Hammermüller S, Kamrath D, Wolf S, Dietrich A, Camilo LM, Beda A, Carvalho AR, Giannella-Neto A. Individualized positive end-expiratory pressure in obese patients during general anaesthesia: a randomized controlled clinical trial using electrical impedance tomography. *BJA: British Journal of Anaesthesia*. 2017 Dec 1;119(6):1194-205.
25. Zorrilla-Vaca A, Grant MC, Urman RD, Frenzl G. Individualised positive end-expiratory pressure in abdominal surgery: a systematic review and meta-analysis. *British journal of anaesthesia*. 2022 Nov 1;129(5):815-25.
26. Kim YJ, Kim BR, Kim HW, Jung JY, Cho HY, Seo JH, Kim WH, Kim HS, Hwangbo S, Yoon HK. Effect of driving pressure-guided positive end-expiratory pressure on postoperative pulmonary complications in patients undergoing laparoscopic or robotic surgery: a randomised controlled trial. *British Journal of Anaesthesia*. 2023 Nov 1;131(5):955-65.
27. Yang G, Zhang P, Li L, Wang J, Jiao P, Wang J, Chu Q. Driving pressure-guided ventilation in obese patients undergoing laparoscopic sleeve gastrectomy: a randomized controlled trial. *Diabetes, Metabolic Syndrome and Obesity*. 2023 Dec 31;15:1515-23.
28. Braun M, Ruscher L, Fuchs A, Kämpfer M, Huber M, Luedi MM, Riva T, Vogt A, Riedel T. Atelectasis in obese patients undergoing laparoscopic bariatric surgery are not increased upon discharge from Post Anesthesia Care Unit. *Frontiers in medicine*. 2023 Aug

29;10:1233609..

29. Ma J, Sun M, Song F, Wang A, Tian X, Wu Y, Wang L, Zhao Q, Liu B, Wang S, Qiu Y. Effect of ultrasound-guided individualized positive end-expiratory pressure on the severity of postoperative atelectasis in elderly patients: a randomized controlled study. *Scientific Reports*. 2024 Nov 15;14(1):28128.
30. Szigetváry C, Szabó GV, Dembrowszky F, Ocskay K, Engh MA, Turan C, Szabó L, Walter A, Kobeissi F, Terebessy T, Hegyi P, Ruszkai Z, Molnár Z. Individualised Positive End-Expiratory Pressure Settings Reduce the Incidence of Postoperative Pulmonary Complications: A Systematic Review and Meta-Analysis. *J Clin Med*. 2024 Nov 11;13(22):6776.
31. Scaramuzzo G, Priani P, Ferrara P, Verri M, Montanaro F, La Rosa R, Cammarota G, Volta CA, Spadaro S. Longitudinal changes of electrical impedance tomography-based best PEEP in obese patients undergoing laparoscopic surgery: a prospective physiological study. *Anaesthesia Critical Care & Pain Medicine*. 2025 Sep 1;44(5):101569.
32. Campos NS, Bluth T, Hemmes SN, Librero J, Pozo N, Ferrando C, Ball L, Mazzinari G, Pelosi P, de Abreu MG, Schultz MJ. Intraoperative positive end-expiratory pressure and postoperative pulmonary complications: a patient-level meta-analysis of three randomised clinical trials. *British journal of anaesthesia*. 2022 Jun 1;128(6):1040-51..
33. Uhlig C, Neto AS, Van Der Woude M, Kiss T, Wittenstein J, Shelley B, Scholes H, Hiesmayr M, Vidal Melo MF, Sances D, Coskunfirat N. Intraoperative mechanical ventilation practice in thoracic surgery patients and its association with postoperative pulmonary complications: results of a multicenter prospective observational study. *BMC anesthesiology*. 2020 Jul 22;20(1):179.
34. Ball L, Almondo C, Pelosi P. Perioperative lung protection: general mechanisms and protective approaches. *Anesthesia & Analgesia*. 2020 Dec 1;131(6):1789-98.

35. Fernandez-Bustamante A, Sprung J, Parker RA, Bartels K, Weingarten TN, Kosour C, Thompson BT, Melo MF. Individualized PEEP to optimise respiratory mechanics during abdominal surgery: a pilot randomised controlled trial. *British journal of anaesthesia*. 2020 Sep 1;125(3):383-92.
36. De Jong A, Wrigge H, Hedenstierna G, Gattinoni L, Chiumello D, Frat JP, Ball L, Schetz M, Pickkers P, Jaber S. How to ventilate obese patients in the ICU. *Intensive care medicine*. 2020 Dec;46(12):2423-35.
37. Loring SH, O'Donnell CR, Behazin N, Malhotra A, Sarge T, Ritz R, Novack V, Talmor D. Esophageal pressures in acute lung injury: do they represent artifact or useful information about transpulmonary pressure, chest wall mechanics, and lung stress?. *Journal of applied physiology*. 2010 Mar;108(3):515-22.
38. Pereira SM, Tucci MR, Morais CC, Simões CM, Tonelotto BF, Pompeo MS, Kay FU, Pelosi P, Vieira JE, Amato MB. Individual positive end-expiratory pressure settings optimize intraoperative mechanical ventilation and reduce postoperative atelectasis. *Anesthesiology*. 2018 Dec 1;129(6):1070-81.
39. Chatfeld AF, Nunez D, Abnet KR, Farhang B, Klick JC, Horn N, Bender SP, Bates JH, Dixon AE, Tharp WG. Effect of Obesity, Pneumoperitoneum, and Body Position on Mechanical Power of Intraoperative 1 Ventilation: An Observational Study 2 3 4 William G. Tharp^{1, 2}, Maegan R. Neilson¹, Max W. Breidenstein^{1, 2}, Ryan G. Harned^{1, 2}, Sydney 5.
40. Cinnella G, Grasso S, Spadaro S, Rauseo M, Mirabella L, Salatto P, De Capraris A, Nappi L, Greco P, Dambrosio M. Effects of recruitment maneuver and positive end-expiratory pressure on respiratory mechanics and transpulmonary pressure during laparoscopic surgery. *Survey of Anesthesiology*. 2013 Dec 1;57(6):282-3.

41. Aldenkortt M, Lysakowski C, Elia N, Brochard L, Tramèr MR. Ventilation strategies in obese patients undergoing surgery: a quantitative systematic review and meta-analysis. *British journal of anaesthesia*. 2012 Oct 1;109(4):493-502.
42. Talab HF, Zabani IA, Abdelrahman HS, Bukhari WL, Mamoun I, Ashour MA, Sadeq BB, El Sayed SI. Intraoperative ventilatory strategies for prevention of pulmonary atelectasis in obese patients undergoing laparoscopic bariatric surgery. *Anesthesia & Analgesia*. 2009 Nov 1;109(5):1511-6.
43. Severgnini P, Selmo G, Lanza C, Chiesa A, Frigerio A, Bacuzzi A, Dionigi G, Novario R, Gregoret C, de Abreu MG, Schultz MJ. Protective Mechanical Ventilation During General Anesthesia for Open Abdominal Surgery Improves Postoperative Pulmonary Function. *Survey of Anesthesiology*. 2014 Feb 1;58(1):19-20.
44. Li, X., Liu, H., Wang, J., Ni, Z. L., Liu, Z. X., Jiao, J. L., Han, Y., & Cao, J. L. (2023). Individualized Positive End-expiratory Pressure on Postoperative Atelectasis in Patients with Obesity: A Randomized Controlled Clinical Trial. *Anesthesiology*, 139(3), 262–273.
45. Boesing C, Schaefer L, Rocco PR, Luecke T, Krebs J. Positive end-expiratory pressure optimisation during general anaesthesia in patients with obesity: a narrative review of respiratory and cardiovascular outcomes. *British Journal of Anaesthesia*. 2025 Oct 16.
46. Feng D, Zhao Y, Wu Z, Li W, Li X, Wan J, Wang F. Ultrasound guided individualized PEEP reduces postoperative atelectasis in elderly patients undergoing laparoscopic radical rectal cancer surgery. *Scientific Reports*. 2025 Nov 12;15(1):39585.
47. Gao L, Zhang B, Qi J, Zhao X, Yan X, Li B, Shen J, Gu T, Yu Q, Luo M, Wang Y. Effects of individualized positive end-expiratory pressure on intraoperative oxygenation and postoperative pulmonary complications in patients requiring pneumoperitoneum with Trendelenburg position: a systematic review and meta-analysis. *International Journal of*

Surgery. 2025 Jan 1;111(1):1386-96.

48. WANG Z, ZHU J, YAO J, ZHANG W, NIE L, SU X, YUAN X, LI C, LIU S, TIAN S. Effect of driving pressure-guided individualized PEEP titration on atelectasis in elderly patients undergoing robot-assisted radical prostatectomy. *Chinese Journal of Anesthesiology*. 2021:1446-50.
49. Vijayakumar, L., Thirunavukkarasu, P. L., & Barathi, A. (2025). Effect of intraoperative positive end-expiratory pressure on post-operative pulmonary complications in overweight patients undergoing elective laparoscopic hernia surgery: A prospective randomised controlled trial. *Journal of minimal access surgery*, 10.4103/jmas.jmas_321_24. Advance online publication.